


INSTRUCTIONS:

Set Weight at Height using the appropriate units of measure.

Read Body Surface Area at arrow.

Please see Precautions and Posology and Method of Administration sections of SmPC for dose modification information.



Body Surface Area (m²)



From the formula of Dubois and Dubois,
Arch Intern Med 1916;17:863


Reconstitution of 3.5 mg VELCADE [®] solution for			
IV injection			
Route of administration	Pack size	Reconstitution volume	Final concentration
Intravenous use only	3.5 mg	3.5 ml	1.0 mg/ml

Reconstitution of 3.5 mg VELCADE [®] solution for			
SC injection			
Route of administration	Pack size	Reconstitution volume	Final concentration
Subcutaneous use only	3.5 mg	1.4 ml	2.5 mg/ml

Subcutaneous or Intravenous use only. Do not give by other routes.

VELCADE[®]
(bortezomib)

Janssen
PHARMACEUTICAL COMPANIES
of Johnson & Johnson





Please include here the abbreviated
SmPC due to your local regulations



Method of administration



IV administration:

The reconstituted solution is administered as a 3-5 second bolus intravenous injection through a peripheral or central intravenous catheter followed by a flush with sodium chloride 9 mg/ml (0.9%) solution for injection.

SC administration:

The reconstituted solution is administered subcutaneously in the thighs (right or left, proximal and distal sites) or abdomen (right or left, upper or lower quadrant). Injection sites could be rotated for successive injections.

Example

Body surface m ²	Total dose required (in mg) with 1.3mg/m ²	Applied volume with IV use (in ml)	Applied volume with SC use (in ml)
1.5	1.95	1.95	0.78
1.6	2.08	2.08	0.83
1.7	2.21	2.21	0.88
1.8	2.34	2.34	0.94
1.9	2.47	2.47	0.99
2.0	2.60	2.60	1.04
2.1	2.73	2.73	1.09