

VFEND® (voriconazole)
oral/intravenous

**Healthcare Professional
Question & Answer Brochure**

CONTENTS

1. What is the purpose of this brochure?.....	3
2. What is VFEND?	3
3. What should I know about phototoxicity and skin squamous cell carcinoma risk associated with VFEND?	4
4. What should I know about patient management to minimise the risk of photosensitivity and squamous cell carcinoma with VFEND?.....	4
5. What do I need to know about the hepatic risk associated with VFEND?	5
6. What are the knowledge and the recommendations regarding patients with hepatic impairment?	5
7. What should I know about safety monitoring to minimise the hepatotoxicity risk of VFEND?.....	5
8. What tools are available to help me for the monitoring of my patients?	6
9. What should I discuss with my patient?.....	7
10. Where can I obtain further information?.....	7
11. How do I report Adverse Reactions/Events?	8

1- WHAT IS THE PURPOSE OF THIS BROCHURE?

These questions and answers (Q&As) are provided by Pfizer Inc. (Pfizer) for prescribers and other healthcare professionals (HCPs) involved in the treatment of patients with VFEND.

This document will enable you to:

- Understand what VFEND is used for and how it should be used
- Be aware of important identified risks of phototoxicity, squamous cell carcinoma (SCC) of the skin and hepatic toxicity adverse reactions of VFEND and how they should be mitigated and managed
- Understand what other tools are available to communicate and remind patients of these risks
- Provide important safety information to patients

Please also familiarise yourself with the complete Summary of Product Characteristics (SmPC), which can be downloaded via the product website, http://www.ema.europa.eu/ema/index.jsp?curl=pages/medicines/human/medicines/000387/human_med_001135.jsp&mid=WC0b01ac058001d124 before prescribing or dispensing VFEND. Please contact Pfizer's Medical Information office at: Telephone: +30 210 6785800, email: eumedinfo@pfizer.com.

2- What is VFEND?

VFEND is a broad spectrum, triazole antifungal agent and is indicated in adults and children aged 2 years and above as follows:

Treatment of invasive aspergillosis.

Treatment of candidemia in non-neutropenic patients.

Treatment of fluconazole-resistant serious invasive *Candida* infections (including *C. krusei*).

Treatment of serious fungal infections caused by *Scedosporium* spp. and *Fusarium* spp.

VFEND should be administered primarily to patients with progressive, possibly life-threatening infections.

Prophylaxis of invasive fungal infections in high risk allogeneic hematopoietic stem cell transplant (HSCT) recipients.

3- What should I know about phototoxicity and skin squamous cell carcinoma risk associated with VFEND?

VFEND has been associated with phototoxicity reactions.

Squamous cell carcinoma of the skin has also been reported in patients receiving VFEND, some of whom have reported prior phototoxic reactions.

4- What should I know about patient management to minimise the risk of phototoxicity and squamous cell carcinoma with VFEND?

All patients, including children, and their parents or caregivers, should be educated about avoiding intense or prolonged exposure to direct sunlight during VFEND treatment and using measures such as protective clothing and sufficient sunscreen with high sun protection factor (SPF).

Patients should be asked to inform you immediately of the occurrence of sunburn or severe skin reaction following exposure to light or sun.

If phototoxic reactions occur, multidisciplinary advice (e.g. a consultation with a dermatologist) should be sought for the patient. VFEND discontinuation should be considered.

Dermatological evaluation should be performed on a systematic and regular basis, whenever VFEND is continued despite the occurrence of phototoxicity-related lesions to allow early detection and management of pre-malignant lesions. VFEND treatment should be discontinued if pre-malignant skin lesions or squamous cell carcinoma are identified.

Squamous cell carcinoma of the skin has been reported in relation with long-term VFEND therapy. Treatment duration with VFEND should be as short as possible and long-term treatment (greater than 6 months) should be considered only if the benefits outweigh the potential risks.

For prophylaxis use, dose adjustments are not recommended in the case of lack of efficacy or treatment-related adverse events. In the case of treatment-related adverse events, discontinuation of voriconazole and use of alternative antifungal agents must be considered.

5- What do I need to know about the hepatic risk associated with VFEND?

VFEND has been associated with hepatic toxicity. In clinical trials, there have been uncommon cases of serious hepatic reactions during treatment with VFEND (including clinical hepatitis, cholestasis, and fulminant hepatic failure including fatalities).

Instances of hepatic reactions were noted to occur primarily in patients with serious underlying medical conditions (predominantly haematological malignancy).

Transient hepatic reactions, including hepatitis and jaundice, have occurred among patients with no other identifiable risk factors.

Liver dysfunction has usually been reversible on discontinuation of therapy.

6- What are the knowledge and the recommendations regarding patients with hepatic impairment?

There are limited data on the safety of VFEND in patients with abnormal liver function tests (LFTs) (aspartate transaminase [AST], alanine transaminase [ALT], alkaline phosphatase [AP], or total bilirubin >5 times the upper limit of normal [ULN]).

Patients with hepatic impairment must be carefully monitored for drug toxicity. In patients with severe hepatic impairment, VFEND must only be used if the benefit outweighs the potential risk.

In patients with mild to moderate hepatic cirrhosis (Child-Pugh A and B) receiving [DRUG], it is recommended that the standard loading dose regimens be used but that the maintenance dose be halved. VFEND has not been studied in patients with severe chronic hepatic cirrhosis (Child-Pugh C).

7- What should I know about safety monitoring to minimise the hepatotoxicity risk of VFEND?

Both children and adult patients receiving VFEND must be carefully monitored for hepatic toxicity.

Clinical management should include laboratory evaluation of hepatic function (specifically AST and ALT) **at the initiation of treatment with VFEND and at least weekly for the first month of treatment.**

Treatment should be as short as possible. However, if based on the benefit-risk assessment the treatment is continued, and if there are no changes in the LFTs, monitoring frequency can be reduced to monthly.

If the LFTs become markedly elevated, VFEND use should be discontinued, unless medical judgment of the risk-benefit of the treatment justifies continued use.

For prophylaxis use, dose adjustments are not recommended in the case of lack of efficacy or treatment-related adverse events. In the case of treatment-related adverse events, discontinuation of voriconazole and use of alternative antifungal agents must be considered.

8- What tools are available to help me for the monitoring of my patients?

The HCP Checklist

The HCP Checklist is a recommended tool. It is designed to help you evaluate and discuss the risks of phototoxicity, SCC of the skin and hepatic toxicity with your patients before prescribing VFEND. This will remind you to closely monitor patients who develop phototoxicity and to refer them for regular dermatological consultation to minimize the risk of developing SCC of the skin, as well as to monitor liver function at the initiation of, and on a regular basis during VFEND treatment.

The completed checklist can be included within the patient chart to document that the patient has been informed of these risks. If other members of your team, such as junior doctors and specialist nurses are involved in treating patients with severe fungal infections, the checklist is a useful educational aid.

The Patient Alert Card

The Patient Alert Card is a folding card, which helps to remind patients about the need for dermatological evaluations on a regular basis (if phototoxic reactions occur). It also urges the patient to report phototoxic symptoms that increase the risk of SCC of the skin.

Additionally, it reminds patients:

- To avoid intense or prolonged exposure to sunlight
- To use protective clothing and sufficient sunscreen with high sun protective factor (SPF)
- To inform their doctor if they develop sunburn or severe skin reactions

You are encouraged to fill in your contact details on the Patient Alert Card and give it to each patient receiving VFEND treatment. Patients should be encouraged to carry this card during their daily activities.

If you need additional copies of the HCP Checklist or the Patient Alert Card, please contact Pfizer Medical Information at Telephone: +30 210 6785800, email: eumedinfo@pfizer.com or download these materials at http://www.ema.europa.eu/ema/index.jsp?curl=pages/medicines/human/medicines/000387/human_med_001135.jsp&mid=WC0b01ac058001d124.

9- WHAT SHOULD I DISCUSS WITH MY PATIENT?

Your role in educating patients about their treatment and its potential adverse effects is very important. You will need to inform patients about:

- Important phototoxicity, SCC of the skin, and hepatic risks associated with VFEND
- The need for dermatological evaluation in case of phototoxicity and regular follow-up afterwards
- The need for patients (including children) to avoid intense or prolonged exposure to direct sunlight during VFEND treatment and to use measures such as protective clothing and sufficient sunscreen with high SPF
- The need for patients to inform you immediately of the occurrence of sunburn or severe skin reaction following exposure to light or sun
- The need for liver function tests on a regular basis
- The need for patients to recognize symptoms and signs of liver toxicity (jaundice, unexplained vomiting, stomach pain, dark urine) and to report to you immediately

You should give the patient a VFEND **Patient Alert Card**, which reinforces the important risk of phototoxicity and skin SCC associated with VFEND treatment, and advise the patient to carry this card during their daily activities.

You should also remind the patient about this important safety information regularly during their treatment with VFEND.

10- Where can I obtain further information?

For further information please contact Pfizer Medical Information at Telephone: +30 210 6785800, email: eumedinfo@pfizer.com .

11- How do I report Adverse Reactions/Events?

It is important that you promptly report any adverse reactions associated with VFEND to assist in fully characterising the product's safety profile.

For any suspected adverse event reactions, please report such events in accordance with the national spontaneous reporting system to the Medicines Authority Post-Licensing Directorate, 203, Level 3, Rue D'Argens, Gzira GZR 1368, Malta, or at: <http://www.medicinesauthority.gov.mt/adrportal> or by contacting Pfizer at Telephone: +30 210 6785800, 24h line: +30 210 6785808 or to the local representative: V.J. Salomone Pharma Ltd. Upper Cross Road, Marsa MRS1542, Malta, Tel: +356 21 220 174.

For further information please contact Pfizer Medical Information at Telephone: +30 210 6785800, email: eumedinfo@pfizer.com