

Package leaflet: Information for the patient

Sasah 50 mg orodispersible film sildenafil

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Sasah is and what it is used for
2. What you need to know before you take Sasah
3. How to take Sasah
4. Possible side effects
5. How to store Sasah
6. Contents of the pack and other information

1. What Sasah is and what it is used for

Sasah contains the active substance sildenafil which belongs to a group of medicines called phosphodiesterase type 5 (PDE5) inhibitors. It works by helping to relax the blood vessels in your penis, allowing blood to flow into your penis when you get sexually excited. Sasah will only help you to get an erection if you are sexually stimulated.

Sasah is a treatment for adult men with erectile dysfunction, sometimes known as impotence. This is when a man cannot get, or keep a hard, erect penis suitable for sexual activity.

2. What you need to know before you take Sasah

Do not take Sasah

- If you are allergic to sildenafil or any of the other ingredients of this medicine (listed in section 6).
- If you are taking medicines called nitrates, as the combination may lead to a dangerous fall in your blood pressure. Tell your doctor if you are taking any of these medicines which are often given for relief of angina pectoris (or “chest pain”). If you are not certain, ask your doctor or pharmacist.

- If you are using any of the medicines known as nitric oxide donors such as amyl nitrite (“poppers”), as the combination may also lead to a dangerous fall in your blood pressure.
- If you are taking riociguat. This drug is used to treat pulmonary arterial hypertension (i.e., high blood pressure in the lungs) and chronic thromboembolic pulmonary hypertension (i.e., high blood pressure in the lungs secondary to blood clots). PDE5 inhibitors, such as sildenafil have been shown to increase the hypotensive effects of this medicine. If you are taking riociguat or are unsure tell your doctor.
- If you have a severe heart or liver problem.
- If you have recently had a stroke or a heart attack, or if you have low blood pressure.
- If you have certain rare inherited eye diseases (such as *retinitis pigmentosa*).
- If you have ever had loss of vision due to non-arteritic anterior ischaemic optic neuropathy (NAION).

Warnings and precautions

Talk to your doctor, pharmacist or nurse before taking Sasah

- If you have sickle cell anaemia (an abnormality of red blood cells), leukaemia (cancer of blood cells), multiple myeloma (cancer of bone marrow).
- If you have a deformity of your penis or Peyronie’s Disease.
- If you have problems with your heart. Your doctor should carefully check whether your heart can take the additional strain of having sex.
- If you currently have a stomach ulcer, or a bleeding problems (such as haemophilia).
- If you experience sudden decrease or loss of vision, stop taking Sasah and contact your doctor immediately.

You should not use Sasah with any other oral or local treatments for erectile dysfunction.

You should not use Sasah with treatments for pulmonary arterial hypertension (PAH) containing sildenafil or any other PDE5 inhibitors.

You should not take Sasah if you do not have erectile dysfunction. You should not take Sasah if you are a woman.

Special considerations for patients with kidney or liver problems

You should tell your doctor if you have kidney or liver problems. Your doctor may decide on a lower dose for you.

Children and adolescents

Sasah should not be given to individuals under the age of 18.

Other medicines and Sasah

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Sasah may interfere with some medicines, especially those used to treat chest pain. In the event of a medical emergency, you should tell your doctor, pharmacist or nurse that you have taken Sasah and when you did. Do not take Sasah with other medicines unless your doctor tells you that you can.

You should not take Sasah if you are taking medicines called nitrates, as the combination of these medicines may lead to a dangerous fall in your blood pressure. Always tell your doctor, pharmacist or nurse if you are taking any of these medicines that are often used for the relief of angina pectoris (or “chest pain”).

You should not take Sasah if you are using any of the medicines known as nitric oxide donors such as amyl nitrite (“poppers”) as the combination may also lead to a dangerous fall in your blood pressure.

Tell your doctor or pharmacist if you are already taking riociguat.

If you are taking medicines known as protease inhibitors, such as for the treatment of HIV, your doctor may reconsider your treatment with sildenafil.

Some patients who take alpha-blocker therapy for the treatment of high blood pressure or prostate enlargement may experience dizziness or light-headedness, which may be caused by low blood pressure upon sitting or standing up quickly. Certain patients have experienced these symptoms when taking sildenafil with alpha-blockers. This is most likely to happen within 4 hours after taking sildenafil. To reduce the chance that these symptoms might happen, you should be on a regular daily dose of your alpha-blocker before you start sildenafil. Your doctor may reconsider your treatment with sildenafil.

Tell your doctor or pharmacist if you are taking, have recently taken or might take medicines containing sacubitril/valsartan, used to treat heart failure.

Sasah with food and drink and alcohol

Sasah can be taken with or without food. However, you may find that Sasah takes longer to start working if you take it with a heavy meal.

Drinking alcohol can temporarily impair your ability to get an erection. To get the maximum benefit from your medicine, you are advised not to drink excessive amounts of alcohol before taking sildenafil.

Pregnancy, breast-feeding and fertility

Sasah is not indicated for use by women.

Driving and using machines

Sasah can cause dizziness and can affect vision. You should be aware of how you react to Sasah before you drive or use machinery.

Contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per film that is to say essentially ‘sodium free’.

This medicine contains 18 mg of propylene glycol (E 1520) in each film.

3. How to take Sasah

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure. The recommended starting dose is 50 mg.

You should not take Sasah more than once a day.

Do not take Sasah orodispersible films in combination with sildenafil tablets.

You should take Sasah about one hour before you plan to have sex. Sasah orodispersible films are for oral use and are taken without water. If desired you may also take Sasah with water.

If you feel that the effect of Sasah is too strong or too weak, talk to your doctor or pharmacist.

Sasah will only help you to get an erection if you are sexually stimulated. The amount of time Sasah takes to work varies from person to person, but it normally takes between half an hour and one hour. You may find that Sasah takes longer to work if you take it with a heavy meal.

If Sasah does not help you to get an erection, or if your erection does not last long enough for you to complete sexual intercourse you should tell your doctor.

If you take more Sasah than you should:

You may experience an increase in side effects and their severity. Doses above 100 mg do not increase the efficacy.

You should not take more Sasah than your doctor tells you to.

Contact your doctor if you take more Sasah than you should.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

How to administer Sasah orodispersible films


- Sildenafil (Sildenafil citrate) is available as 50 mg orodispersible film that dissolves in the mouth.
- Open the Sasah orodispersible film sachet with dry hands.
- The orodispersible film should be placed on your tongue, where it dissolves and can be

swallowed with the saliva.

- The orodispersible film can be used in situations in which liquids are not available, or to avoid the nausea and vomiting that may accompany the ingestion of film with liquids.

Instructions for use

Important: Do not handle the orodispersible film with wet hands!



Directions for use:

Step 1. Hold the sachet with backside facing you. Locate the arrow on the side, which is at the top left corner of the sachet and hold it with the side facing up. The sachet is partially sealed on this side with two exposed flaps.

Step 2. Hold the two exposed flaps of the sachet at the arrow mark. With a gentle peeling motion, start pulling apart both flaps of the sachet in opposite directions and thereby separating the two layers of the sachet.

Step 3. Continue the peeling motion until the two layers of the sachet are now almost separated. The orodispersible film is now visible inside the sachet. There is no need to completely pull apart both the layers of the sachet to access the orodispersible film.

Step 4, 5 and 6. Carefully remove the orodispersible film from the sachet and put the orodispersible film on top of the tongue. It will itself get dissolve within seconds.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. The side effects reported in association with the use of sildenafil are usually mild to moderate and of a short duration.

If you experience any of the following serious side effects stop taking Sasah and seek medical help immediately:

- An allergic reaction - this occurs **uncommonly** (may affect up to 1 in 100 people) Symptoms include sudden wheeziness, difficulty in breathing or dizziness, swelling of the eyelids, face, lips or throat.
- Chest pains - this occurs uncommonly
If this occurs during or after intercourse

- Get in a semi-sitting position and try to relax.
 - **Do not use nitrates** to treat your chest pain.
- Prolonged and sometimes painful erections - this occurs **rarely** (may affect up to 1 in 1,000 people)
If you have an erection which lasts for more than 4 hours, you should contact a doctor immediately.
 - A sudden decrease or loss of vision - this occurs **rarely**
 - Serious skin reactions - this occurs **rarely**
Symptoms may include severe peeling and swelling of the skin, blistering of the mouth, genitals and around the eyes, fever.
 - Seizures or fits - this occurs **rarely**

Other side effects:

Very common (may affect more than 1 in 10 people): headache.

Common (may affect up to 1 in 10 people): nausea, facial flushing, hot flush (symptoms include a sudden feeling of heat in your upper body), indigestion, colour tinge to vision, blurred vision, visual disturbance, stuffy nose and dizziness.

Uncommon (may affect up to 1 in 100 people): vomiting, skin rash, eye irritation, bloodshot eyes /red eyes, eye pain, seeing flashes of light, visual brightness, light sensitivity, watery eyes, pounding heartbeat, rapid heartbeat, high blood pressure, low blood pressure, muscle pain, feeling sleepy, reduced sense of touch, vertigo, ringing in the ears, dry mouth, blocked or stuffy sinuses, inflammation of the lining of the nose (symptoms include runny nose, sneezing and stuffy nose), upper abdominal pain, gastro-oesophageal reflux disease (symptoms include heartburn), presence of blood in urine, pain in the arms or legs, nosebleed, feeling hot and feeling tired.

Rare (may affect up to 1 in 1,000 people): fainting, stroke, heart attack, irregular heartbeat, temporary decreased blood flow to parts of the brain, feeling of tightening of the throat, numb mouth, bleeding at the back of the eye, double vision, reduced sharpness of vision, abnormal sensation in the eye, swelling of the eye or eyelid, small particles or spots in your vision, seeing halos around lights, dilation of the pupil of the eye, discolouration of the white of the eye, penile bleeding, presence of blood in semen, dry nose, swelling of the inside of the nose, feeling irritable and sudden decrease or loss of hearing.

From post-marketing experience cases of unstable angina (a heart condition) and sudden death have been reported rarely. Of note, most, but not all, of the men who experienced these side effects had heart problems before taking this medicine. It is not possible to determine whether these events were directly related to sildenafil.

Reporting of side effects

If you get any side effects, talk to your doctor pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via: ADR Reporting Website:

<https://medicinesauthority.gov.mt/adrportal>. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Sasah

Keep this medicine out of the sight and reach of children.

This medicinal product does not require any special storage conditions.

Do not use this medicine after the expiry date, which is stated on the carton/sachet after "EXP". The expiry date refers to the last day of that month.

Do not remove the orodispersible film from the outer aluminium sachet until you are ready to take the medicine inside. Do not use the medicine if you notice that the aluminium sachet is damaged.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Sasah contains

- The active substance is sildenafil. Each orodispersible film contains sildenafil citrate equivalent to 50 mg of sildenafil.
- The other ingredients are:

Hypromellose 50 cps (E 464)
Propylene Glycol (E 1520)
Povidone (PVP K30)
Sodium Carbonate Anhydrous Polysorbate 80
Poly (Vinyl alcohol) Sucralose
Hypromellose 15 cps (E 464)
Sodium Laurilsulfate
Neotame
Bitterness Masker (B.T.M.)
Raspberry Flavor
Mint Oil Partly Dementholised
FD & C Blue 1 color (E 133)

What Sasah looks like and contents of the pack

Sasah 50 mg is blue, rectangular, opaque, non-sticky orodispersible film.

Sasah 50 mg orodispersible film is available in a triple laminate foil sachet containing single orodispersible film.

Pack of 2, 4, 6, 8 or 10 orodispersible films in a carton.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

Marketing Authorisation Holder

Novatin Limited
230, Second floor
Eucharistic Congress Road
Mosta, MST 9039
Malta

Manufacturer

Laboratori Fundació Dau
Pol. Ind. Consorci Zona Franca. c/C, 12-14
Barcelona, 08018 Spain

This medicine is authorised in the Member States of the European Economic Area under the following names:

Name of the Member State	Name of the medicine
Czechia	Sasah
Croatia	Sasah 50 mg raspadljivi film za usta
Malta	Sasah 50 mg orodispersible film

This leaflet was last revised in 10/2022