

Package leaflet: Information for the user

Solpadeine soluble tablets

Paracetamol 500mg, codeine phosphate hemihydrate 8mg and caffeine 30mg

Read all of this leaflet carefully before you start using this medicine because it contains important information for you.

Always use this medicine exactly as described in this leaflet or as your doctor or pharmacist have told you.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.
- You must talk to a doctor if you do not feel better or if you feel worse after 3 days.

What is in this leaflet

1. What this medicine is and what it is used for
2. What you need to know before you take this medicine
3. How to take this medicine
4. Possible side effects
5. How to store this medicine
6. Contents of the pack and other information

1. What this medicine is and what it is used for

Solpadeine soluble tablets contains the active substances paracetamol 500 mg and codeine phosphate hemihydrate 8 mg , which are both pain-relieving drugs, and caffeine 30 mg, a stimulant which acts to enhance the pain-relieving effects of paracetamol.

Codeine belongs to a group of medicines called opioid analgesics which act to relieve pain. It can be used on its own or in combination with other pain killers such as paracetamol. Paracetamol also helps to reduce temperature when you have a fever. Codeine has some cough suppressant action.

Solpadeine soluble tablets can be used in patients over 12 years of age for the short-term relief of acute, moderate pain that is not relieved by other painkillers such as paracetamol or ibuprofen alone. This medicine is not recommended for use in children aged 12 years to 18 years with compromised respiratory function for the symptomatic treatment of cough and/or cold.

Solpadeine soluble tablets provide effective relief from pain, including:

- Headache, including migraine
- Toothache
- Period pain
- Rheumatic pain
- Backache
- Neuralgia
- Relief of cold and flu symptoms

This product should be used only when clearly necessary.

2. What you need to know before you take this medicine

Do not take this medicine

- if you are allergic to paracetamol, codeine, caffeine or any of the other ingredients of this medicine (listed in section 6);
- if you suffer from acute asthma (if you have suffered from recent asthma attacks);
- in children and adolescents (0-18 years of age) after removal of their tonsils or adenoids due to obstructive sleep apnoea syndrome;
- if you are breastfeeding;
- if you know that you metabolise very rapidly codeine into morphine;
- if you are under 12 years old;
- if you suffer from chronic constipation.

Warnings and precautions

Talk to your doctor or pharmacist before taking this medicine:

- if you suffer from an enlarged prostate (difficulty passing water);
- if you suffer from any bowel disorders;
- if you suffer from any liver or kidney problems;
- if you suffer from heart or blood pressure problems;
- if you have been told by your doctor that you have an intolerance to some sugars;
- if you have recently had gall bladder surgery or a similar operation.
- if you have chronic alcoholism;
- if you have Gilbert's Syndrome (familial non-haemolytic jaundice);
- if you are taking other medicines that can affect the liver;
- if you have glucose-6-phosphate dehydrogenase deficiency;
- if you have haemolytic anaemia;
- if you are dehydrated;
- if you have had poor diet such that the diet causes health problems;
- if you weigh less than 50kg;
- if you are elderly.

They may reduce the dose or increase the time interval between doses of your medicine.

Talk to your doctor or pharmacist before taking Neo-codion:

- if you are or have ever been addicted to opioids, alcohol, prescription medicines, or illegal substances

Taking codeine (an active ingredient of this medicine) regularly for a long time can lead to addiction and misuse, which may result in overdose and/or death.

Do not take this medicine longer than needed. Do not give this medicine to anyone else.

Codeine is transformed to morphine in the liver by an enzyme. Morphine is the substance that produces pain relief. Some people have a variation of this enzyme and this can affect people in different ways. In some people, morphine is not produced or produced in very small quantities, and it will not provide enough pain relief.

Other people are more likely to get serious side effects because a very high amount of morphine is produced. If you notice any of the following side effects, you must stop taking this medicine and seek immediate medical advice: slow or shallow breathing, confusion, sleepiness, small pupils, feeling or being sick, constipation, lack of appetite.

Special warnings about addiction

This product contains codeine. This medicine should only be taken when necessary. Do not take more than the stated dose and do not take every day for more than 3 days unless told to do so by your doctor. Prolonged regular use, except under medical supervision, may lead to addiction and result in withdrawal symptoms, such as restlessness and irritability once the drug is stopped. If you find you need to use this product all the time or think you are suffering from withdrawal symptoms, it is important to consult your doctor or pharmacist.

Children and adolescents

This medicine should not be used for pain relief in children and adolescents after removal of their tonsils or adenoids due to Obstructive Sleep Apnoea Syndrome. Codeine is not recommended in children or adolescents with breathing problems, since the symptoms of morphine toxicity may be worse in these children.

Other medicines and Solpadeine

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, especially:

- medicines that you have bought yourself such as cough/cold remedies and other painkillers. Many of these will contain paracetamol and/or codeine and must not be taken while you are taking Solpadeine soluble tablets;
- if you are taking or have taken within the last 2 weeks medicines prescribed by your doctor for the treatment of depression or anxiety, or medicines known as tranquilisers (including monoamine oxidase inhibitors-MAOIs);
- medicines used to relax you or to help you sleep (such as benzodiazepines);
- medicines for high cholesterol levels or for the treatment of diarrhoea (cholestyramine);
- medicines to stop you feeling sick or vomiting (domperidone or metoclopramide);
- medicines to prevent blood clots (such as warfarin);
- medicines for the treatment of gout (probenecid);
- medicines for eye infections (chloramphenicol);
- anticonvulsants and medicines used to treat epilepsy (such as barbiturates);
- oral contraceptive steroids.
- flucloxacillin (antibiotic), due to a serious risk of blood and fluid abnormality (high anion gap metabolic acidosis) that must have urgent treatment and which may occur particularly in case of severe renal impairment, sepsis (when bacteria and their toxins circulate in the blood leading to organ damage), malnutrition, chronic alcoholism, and if the maximum daily doses of paracetamol are used.

Concomitant use of this medicine and sedative medicines such as benzodiazepines or related drugs increases the risk of drowsiness, difficulties in breathing (respiratory depression), coma and may be life-threatening. Because of this, concomitant use should only be considered when other treatment options are not possible.

However if your doctor does prescribe this medicine together with sedative medicines the dose and duration of concomitant treatment should be limited by your doctor.

Please tell your doctor about all sedative medicines you are taking and follow your doctor's dose recommendation closely. It could be helpful to inform friends or relatives to be aware of the signs and symptoms stated above. Contact your doctor when experiencing such symptoms.

Solpadeine with food, drink and alcohol

Do not drink alcohol when taking this medicine.

Avoid excessive intake of caffeine containing products such as coffee, tea and some canned drinks.

Pregnancy, breast-feeding and fertility

Do not use this medicine while you are breastfeeding, it may be present in breast milk.

Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

This medicine may make you feel drowsy. You should not drive or operate machinery until you know how this medicine affects you.

This medicine contains sodium

This medicine contains 427 mg sodium in each tablet. The maximum recommended daily dose of this medicinal product contains 3416 mg sodium (found in table salt). This is equivalent to 171% of the

adult recommended maximum daily dietary intake for sodium. Talk to your pharmacist or doctor if you are taking Solpadeine soluble tablets and have been advised to have a low salt diet.

This medicine contains 50mg sorbitol (E 420) per tablet. Sorbitol is a source of fructose. If your doctor has told you that you (or your child) have an intolerance to some sugars or if you have been diagnosed with hereditary fructose intolerance (HFI), a rare genetic disorder in which a person cannot break down fructose, talk to your doctor before you (or your child) take or receive this medicine.

3. How to take this medicine

Always take this medicine exactly as described in this leaflet or as your doctor or pharmacist have told you. Check with your doctor or pharmacist if you are not sure.

The recommended dose is:

Adults

1-2 tablets every 4-6 hours. Do not take more than 4 doses in 24 hours, equivalent to 8 tablets. The dose should not be repeated more frequently than every 4 hours. Elderly patients may require a reduced dose or frequency of dosing.

Use in children and adolescents

Adolescents 16 to 18 years of age

1-2 tablets every 6 hours. Do not take more than 8 tablets in 24 hours.

Adolescents 12 to 15 years of age

1 tablet every 6 hours. Do not take more than 4 tablets in 24 hours.

Children aged less than 12 years:

Codeine should not be used in children below the age of 12 years, due to the risk of severe breathing problems.

Method of administration

Dissolve the tablets in at least half a glass of water.

Do not exceed the stated dose.

This medicine should not be taken for more than 3 days. If the pain does not improve after 3 days, talk to your doctor for advice. Prolonged use except under medical supervision may be harmful.

If you take more Solpadeine than you should

Talk to a doctor at once if you take too much of this medicine, even if you feel well. This is because too much paracetamol can cause delayed, serious liver damage.

The symptoms can include: nausea, vomiting, anorexia, pale skin, and abdominal pain, or patients may be asymptomatic.

If you forget to take this medicine

Do not take a double dose to make up for a forgotten dose. You must allow at least 4 hours between doses.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Prolonged regular use, except under medical supervision, may lead to addiction and result in withdrawal symptoms, such as restlessness and irritability once the drug is stopped. If you need to take

the medicine for longer periods of time or you need to take more than the recommended dose or feel irritable and restless when you stop taking the medicine, you may be addicted. In any of these apply to you, it is important that you talk to your doctor.

This medicine may make you feel dizzy or drowsy. You may also feel sick or vomit. These effects may be relieved by lying down. You may also become constipated. You may experience problems with your blood which means that you may bleed or bruise easily (known as thrombocytopenia) or which makes it more likely that you may get an infection (known as agranulocytosis).

If you have previously had gall bladder removal surgery and experience severe abdominal pain, nausea and vomiting, or if you have previously experienced breathing problems with aspirin or other non-steroidal anti-inflammatories and experience a similar reaction with this product, stop taking this medicine and tell your doctor immediately.

Taking any type of pain reliever for headaches too often or for a too long period of time can make them worse. If this situation happens to you talk to your doctor or pharmacist.

Some of the following side effects may occur:

- allergic reactions such as skin rash or itching, sweating, sometimes with breathing problems, or swelling of the lips, tongue, throat or face;
- unusual bleeding or bruising under the skin
- skin rash or peeling of the skin which may be accompanied by mouth ulcers;
- severe skin reactions such as severe blisters and bleeding in the lips, eyes, mouth, nose and genitals and formation of blisters containing pus;
- blood disorders including low platelet count (making you slow to heal and bruise easily) and low white blood cell count (making you more likely to get infections);
- cloudy urine or urinary retention;
- nervousness and tremor;
- hallucinations and nightmares;
- fast or irregular heartbeat;
- blurred vision;
- facial flushing;
- constipation.

These effects go away once you stop taking the medicine. If any of these effects occur, or get serious, or if you notice any other side effect not listed in this leaflet, stop taking this medicine, and tell your doctor or pharmacist immediately.

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via HPRa Pharmacovigilance; Website: www.hpra.ie. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store this medicine

Keep out of the reach and sight of children.

Do not use this medicine after the expiry date which is stated on the label. The expiry date refers to the last day of that month.

Do not store this medicine above 25°C.

Store in original packaging to protect from light and moisture.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Further information

What this medicine contains

- The active substances are paracetamol 500mg, codeine phosphate hemihydrate 8mg and caffeine 30mg.
- The other ingredients are sodium hydrogen carbonate, sorbitol (E420), saccharin sodium, sodium laurilsulfate, anhydrous citric acid, anhydrous sodium carbonate, povidone and dimeticone.

What this medicine looks like and contents of the pack

Solpadeine soluble tablets are flat, white soluble tablets, with bevelled edges, plain on one side, breakline on the other. They are available in packs of 4, 8, 12, 24, 32, 36, 48 or 60 tablets. Not all pack sizes may be marketed.

Marketing Authorisation Holder

Chefaro Ireland DAC
The Sharp Building
Hogan Place
Dublin 2
Ireland

All enquiries should be sent to this address.

Manufacturer

GlaxoSmithKline Dungarvan Limited
Knockbrack, Dungarvan, Co. Waterford
or
Swiss Caps GmbH
Grassingerstraße 9
83043 Bad Aibling
Germany (Aenova Group).

This leaflet was last revised in May 2022