

**Read all of this leaflet carefully because it contains important information for you.**

This medicine is available without prescription. However, you still need to use Anginovag carefully to get the best results from it

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- You must contact a doctor if your symptoms worsen or do not improve after 3 days
- If any of side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist

**In this leaflet:**

1. What ANGINOVAG is and what it is used for
2. Before you use ANGINOVAG
3. How to use ANGINOVAG.
4. Possible side effects.
5. How to store ANGINOVAG
6. Further information

**1. WHAT Anginovag IS AND WHAT IT IS USED FOR:**

Anginovag is an oromucosal spray

Anginovag is supplied in a white polyethylene flask containing a solution and closed by an spray dosing valve.

Anginovag is used to alleviate the symptoms of buccopharyngeal affections (pain, inflammation, etc)

**2. BEFORE YOU USE Anginovag:**

**Do not use Anginovag:**

- If you are allergic (hypersensitive) to active substances or any of the other ingredients of Anginovag.

**Take special care with Anginovag**

Do not spray on the eyes

Contact your doctor if you experience blurred vision or other visual disturbances

**Taking other medicines:**

You can take other medication by oral or injection route.

It is advisable do not use Anginovag with other topical medication

**Using Anginovag with food and drink**

It is possible to use Anginovag after the meals.

It is not advisable to take foods or to drink after to use Anginovag

**Pregnancy and breast-feeding**

Anginovag can be used during pregnancy or breast-feeding.

**Driving and using machines**

Anginovag, does not produce any effect over the ability to drive and use of machines

### **Important information on some of the ingredients of Anginovag**

This medicine contains ethanol and sodium saccharin.

### **3. HOW TO USE Anginovag**

The usual dose is:

The first 3 days, 1-2 applications each 2-3 hours and after 1-2 applications each 6 hours.

#### **Method of administration:**

It is recommended to place the flask in upright position with the cannula pointing towards the place of application, but without introducing it completely in the mouth.

Press the upper part of the cap.

The usual duration of the therapy is one week

#### **If you use more Anginovag than you should:**

If you take more Anginovag than indicated, it is not foreseen any complication.

#### **If you forget to use Anginovag:**

Any problem. You can advance the next administration

#### **If you stop using Anginovag**

It is possible that your symptoms worsen.

### **4. POSSIBLE SIDE EFFECTS**

Like all medicines, **Anginovag** can have side effects.

With not known frequency (cannot be estimated from the available data), blurred vision may occur.

If any of side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist

### **5. HOW TO STORE Anginovag**

Keep out of the reach and sight of children.

Do not use Anginovag after the expiry date which is stated on the label and carton. [Cad./Exp./Pér.: number of month-year] The expiry date refers to the last day of that month.

Store at a temperature under 30°C

Do not use Anginovag if you notice visible signs of deterioration.

## **6. FURTHER INFORMATION**

### **What Anginovag contains**

#### ***Active substances***

Dequalinium chloride 0.100g; Tyrothricin 0.400g; Hydrocortisone acetate 0.060g;  
Lidocaine hydrochloride 0.100 g; beta-Glycyrrhetic acid 0.060 g

#### ***Other ingredients***

Sodium saccharin, Propyleneglycol, Pineapple oil, Ethyl alcohol 96°

### **What Anginovag looks like and contents of the pack**

White polyethylene flask provide with an spray dosing valve, and a blue label.  
Bottles containing 10 mL of solution

### **Marketing authorization Holder and Manufacturer**

FERRER INTERNACIONAL, S.A.  
Gran Vía Carlos III, 94  
08028 – Barcelona (Spain)

**This leaflet was approved in- April 2020.**