

PRADAXA® (DABIGATRAN ETEXILATE)

EDUCATIONAL PACK

For prevention of stroke and systemic embolism in adult patients with nonvalvular atrial fibrillation with one or more of the following risk factors:

- Previous stroke, transient ischaemic attack or systemic embolism (SEE)
- Left ventricular ejection fraction <40%
- Symptomatic heart failure, ≥ New York Heart Association (NYHA) Class 2
- Age ≥75 years
- Age ≥65 years associated with one of the following: diabetes mellitus, coronary artery disease or hypertension^{1,2}

References:

1. Boehringer Ingelheim. Pradaxa® 150mg hard capsules Summary of Product Characteristics.
2. Boehringer Ingelheim. Pradaxa® 110mg hard capsules Summary of Product Characteristics.

Prescribing Information (SPAF – UK) PRADAXA® (dabigatran etexilate)

Capsules containing 110 mg or 150 mg dabigatran etexilate (as mesilate) **Action:** Direct thrombin inhibitor. **Indication:** Prevention of stroke and systemic embolism in adult patients with nonvalvular atrial fibrillation with one or more of the following risk factors: Previous stroke, transient ischaemic attack, or systemic embolism (SEE); Left ventricular ejection fraction < 40 %; Symptomatic heart failure, ≥ New York Heart Association (NYHA) Class 2; Age ≥ 75 years; Age ≥ 65 years associated with one of the following: diabetes mellitus, coronary artery disease, or hypertension. **Dose and Administration:** Renal function should be assessed by calculating CrCL prior to initiation to exclude patients with severe renal impairment (CrCL < 30 mL/min). Recommended daily dose 300 mg taken as one 150 mg capsule twice daily. Therapy should be continued long term. In case of interability to dabigatran, patients should be instructed to immediately consult their doctor. **Elderly:** Aged ≥ 80 years 220 mg taken as one 110 mg capsule twice daily; 75 – 80 years consider 220 mg taken as one 110 mg capsule twice daily. As renal impairment may be frequent in the elderly (> 75 years), assess renal function by calculating CrCL prior to initiation to exclude patients with severe renal impairment (CrCL < 30 mL/min). Renal function should also be assessed at least once a year or more frequently as needed in certain clinical situations when it is suspected that the renal function could decline or deteriorate. Patients with an increased risk of bleeding, closely monitor clinically looking for signs of bleeding or anaemia. Dose adjustment should be decided at the discretion of the physician, following assessment of the potential benefit and risk to an individual patient. A coagulation test may help identify increased risk patients. Patients with gastritis, oesophagitis, or gastroesophageal reflux consider 220 mg taken as one 110 mg capsule twice daily due to the elevated risk of major gastro-intestinal bleeding. Renal impairment: contraindicated in severe renal impairment (CrCL < 30 mL/min), patients with renal impairment and a high risk of bleeding consider 220 mg taken as one 110 mg capsule twice daily. Close clinical surveillance is recommended in patients with renal impairment. As above assess renal function prior to initiation to exclude patients with severe renal impairment and assess renal function at least once a year or more frequently as needed. Concomitant verapamil 220 mg taken as one 110 mg capsule twice daily: Pradaxa and verapamil should be taken at the same time. No dose adjustment required but close clinical surveillance in patients < 50 kg. Not recommended if liver enzymes > 2 Upper Limit of Normal (ULN). If switching from Pradaxa to parenteral anticoagulant wait 12 hours after the last dose of Pradaxa; if switching from parenteral anticoagulants to Pradaxa then Pradaxa should be given 0-2 hours prior to the time that the next dose of the alternate therapy would be due, or at the time of discontinuation in case of continuous treatment; if switching from Pradaxa to VKA adjust the starting time of the VKA based on CrCL; if switching from VKA to Pradaxa stop VKA and give Pradaxa once INR < 2.0. Cardioversion patients can stay on Pradaxa whilst being cardioverted. No relevant use of Pradaxa in the paediatric population in the indication. Pradaxa should be swallowed whole with water, with or without food. Patients should be instructed not to open the capsule as this may increase the risk of bleeding. **Contraindications:** Hypersensitivity to any component; severe renal impairment (CrCL < 30 mL/min); active clinically significant bleeding; lesion or condition at significant risk of major bleeding such as current or recent gastrointestinal ulceration, presence of malignant neoplasms at high risk of bleeding, recent brain or spinal injury, recent brain, spinal or ophthalmic surgery, recent intracranial haemorrhage, known or suspected cephalopod venous, arteriovenous malformations, vascular aneurysms or major intrasplenic or intracerebral vascular abnormalities; concomitant treatment with any other anticoagulants e.g. unfractionated heparin (UFH), low molecular weight heparins (enoxaparin, dalteparin etc), heparin derivatives (fondaparinux etc), oral anticoagulants (warfarin, rivaroxaban, apixaban etc) except under the circumstances of switching therapy to or from Pradaxa or when UFH is given at doses necessary to maintain an open central venous or arterial catheter; hepatic impairment or liver disease expected to have any impact on survival; concomitant systemic ketoconazole, cyclosporine, itraconazole, tacrolimus, dioneosone; prosthetic heart valves requiring anticoagulant treatment. **Warnings and Precautions:** Not recommended if liver enzymes > 2 ULN. Haemorrhagic risk. Close clinical surveillance (signs of bleeding or anaemia) is recommended throughout the treatment period, especially when haemorrhagic risk is increased or risk factors combined. Factors which may increase

haemorrhagic risk: age ≥ 75 years; moderate renal impairment (CrCL 30 – 50 mL/min); P-glycoprotein inhibitor co-medication; body weight < 50 kg; acetylsalicylic acid (aspirin); NSAID; clopidogrel; selective serotonin re-uptake inhibitors (SSRIs) or selective serotonin norepinephrine re-uptake inhibitors (SNRIs); other drugs which may impair haemostasis; diseases/procedures associated with a risk of bleeding such as coagulation disorders, thrombocytopenia or functional platelet defects; recent biopsy, major trauma, bacterial endocarditis, oesophagitis, gastritis or gastroesophageal reflux. The measurement of dabigatran related anticoagulation may be helpful to avoid excessive high exposure to dabigatran in the presence of additional risk factors. Patients who develop acute renal failure must discontinue Pradaxa. If severe bleeding occurs, discontinue treatment and investigate the source of the bleeding. Avoid or use with caution medicinal products which may increase the risk of haemorrhage. The use of fibrinolytic medicinal products for the treatment of acute ischaemic stroke may be considered if the patient presents with a T, ECT or aPTT not exceeding the ULN according to the local reference range. Avoid concomitant administration with P-gp inducers. Patients on dabigatran etexilate who undergo surgery or invasive procedures are at increased risk for bleeding therefore surgical interventions may require the temporary discontinuation of dabigatran etexilate; prescribers should consult the Summary of Product Characteristics for further information. Procedures such as spinal anaesthesia may require complete haemostatic function. The risk of spinal or epidural haematoma may be increased in cases of traumatic or repeated puncture and by the prolonged use of epidural catheters. After removal of a catheter, an interval of at least 2 hours should elapse before the administration of the first dose of dabigatran etexilate; these patients require frequent observation for neurological signs and symptoms of spinal or epidural haematoma. Treat with caution patients at high surgical mortality risk and with intrinsic risk factors for thrombotic events: Myocardial infarction. Contains Sunset Yellow (E110) which may cause allergic reactions. **Interactions:** Anticoagulants and antiplatelet aggregation medicinal products; Strong P-gp inhibitors e.g. amiodarone, quinidine, verapamil, clarithromycin co-administration (close clinical surveillance); verapamil co-administration – reduce Pradaxa dose to 220 mg (see above); not recommended for concomitant treatment posaconazole, protease inhibitors including ritonavir and its combinations with other protease inhibitors; avoid with P-gp inducers e.g. rifampicin, St John's wort, carbamazepine, phenytoin, SSRIs or SNRIs. Dabigatran etexilate and dabigatran are not metabolised by cytochrome CYP450 system, therefore related medicinal product interactions not expected. Pantoprazole and other proton-pump inhibitors (PPI) were co-administered with Pradaxa in clinical trials and concomitant PPI treatment did not appear to reduce the efficacy of Pradaxa. Rantidolone administration together with Pradaxa had no clinically relevant effect on the extent of absorption of dabigatran. **Fertility, pregnancy and lactation:** Avoid pregnancy during treatment. Do not use in pregnancy unless clearly necessary. Discontinue breast-feeding during treatment. **Undesirable effects:** Most commonly reported adverse reactions are bleedings occurring in total in approximately 16.5 % in patients with atrial fibrillation treated for the prevention of stroke and SEE. Common (≥ 1/100 to <1/10): anaemia, epistaxis; gastrointestinal haemorrhage; abdominal pain; diarrhoea, dyspepsia, nausea; hepatic function abnormal/ liver function test abnormal; genitourinary haemorrhage. Prescribers should consult the Summary of Product Characteristics for further information on side effects. **Pack sizes and NHS price:** 110 mg 60 capsules £65.90 150 mg 60 capsules £65.90 **Legal category POM MA numbers:** 110 mg EU/1/08/442/007 (60 capsules) 150 mg EU/1/08/442/011 (60 capsules) **Marketing Authorisation Holder:** Boehringer Ingelheim International GmbH, Binger Str. 173, D-55216 Ingelheim am Rhein, Germany. Prescribers should consult the Summary of Product Characteristics for full prescribing information. **Prepared in February 2013.**

Adverse events should be reported. Reporting forms and information can be found at www.mhra.gov.uk/ Yellowcard. Adverse events should also be reported to Boehringer Ingelheim Drug Safety on 0900 328 1627 (frampton).

Boehringer Ingelheim is pleased to provide you with this Educational Pack which has been developed to give practical and relevant information on the appropriate use of Pradaxa®. The pack includes:

- Pradaxa® 150mg – Summary of Product Characteristics
- Pradaxa® 110mg – Summary of Product Characteristics
- Prescriber Guide – this addresses recommendations for the use of Pradaxa® in order to minimise the risk of bleeding
- Patient Alert Card

To order additional copies of the Patient Alert Card please go to: www.pradaxa.co.uk/SPAFeducationalpack

You can also order or download this Educational Pack.



Date of preparation: February 2013 Job code: UK/DBG-121533(1) 