



Asthma

A Parent's Guide



Peter Greally
MD, FRCPI, FRCPC, DCH

AstraZeneca 
Respiratory





Page No. Contents

- | | |
|----|--|
| 1 | Childhood asthma
What is asthma? |
| 2 | Symptoms
Causes
How is asthma diagnosed?
Managing Asthma |
| 3 | Trigger factors |
| 6 | Medical treatment of asthma |
| 8 | Complimentary Therapy
Inhaler devices and your child |
| 9 | Tips for giving your child their inhaler
Nebulisers
Asthma & Rhinitis |
| 10 | Knowing when your child's asthma becomes worse?
The future for asthma
Asthma attack – how can you help? |

This booklet is part of a series on respiratory disease. It endeavours to help parents gain a better understanding regarding their child's asthma. It is not intended as a textbook on asthma. Always follow your doctor's advice and contact them for any further information.

AstraZeneca 
Respiratory





Childhood Asthma


The rate of rise in asthma appears to have reached a plateau. Ireland has one of the highest rates of asthma in Europe and is currently ranked fourth highest in the world for asthma prevalence. As many as 470,000 Irish people have asthma and 1 in 5 Irish children will have asthmatic symptoms at some point in childhood. [Source: Asthma Society of Ireland].

It can be frightening to discover that your child has asthma, however, it is reassuring to know that asthma can be well controlled with the appropriate medications. We hope that this booklet will answer questions you might have.

What is asthma?

The lungs consist of a series of tiny branching tubes called airways, which carry air in and out of little air sacs (alveoli), exchanging oxygen for carbon dioxide. The airways (or bronchial tubes) have muscles within their walls and a lining layer (see diagram). In asthma, the airways become more narrow than normal so it is more difficult for air to pass into the lungs and for used air to be exhaled. The sensation is like breathing in and out through a straw.

- The muscle layer tightens around the airways
- The lining layer becomes swollen and inflamed
- The airways make more mucus than usual





Symptoms

1. Cough: after exercise and/or at night.

A cough may be the only symptom of asthma.

2. Wheeze: whistling sound heard most often when breathing out.
3. Shortness of breath and complaints of tightness in the chest.
4. Lack of stamina during sport/exertion.
Complaints of tiredness.
5. In severe attacks: laboured breathing, blue discolouration of the lips/tongue.

Causes

The exact cause of asthma is as yet unknown. We do know however, that asthma can run in families along with other allergic conditions such as hay fever, allergic conjunctivitis (itchy eyes) and eczema. Outdoor pollution, although not proven to be a cause of asthma, can make symptoms worse. We also know that smoking during pregnancy and passive smoking can increase symptoms in babies and children.

How is asthma diagnosed?

Asthma can be difficult to diagnose in young children.

A diagnosis is usually made on a pattern of symptoms over a period of time. A breathing test (peak flow or pulmonary function test) may be carried out on older children, or sometimes a bronchial challenge test is required. This will usually be performed by a technician in a pulmonary function laboratory.

Unfortunately there is no simple blood, breathing or x-ray test that can reliably diagnose asthma. Often small children are given a trial of medication when asthma is suspected.

It may take several visits to your doctor before the diagnosis is made.

Managing asthma

Asthma cannot be cured but symptoms can be successfully controlled by avoidance of known triggers and by taking medication. It is very important that your child's controller medications are taken regularly.





Trigger Factors

In asthma, the inflamed airways are more irritable than normal. As a result they are quick to narrow in response to various stimuli:



Virus infections: head colds, sore throats and flu



Fumes: tobacco smoke, fire smoke, paint, perfumes and air pollution



Exercise, excitement, upset and worry



Allergies e.g. house dust mite



Pets and pollens



Weather: cold air, damp conditions and fog





House dust mites

These are microscopic creatures that feed off human skin scales and thrive very well in warm humid atmospheres. Some simple measures can help reduce the number of dust mites:



1. Bed linen should be washed weekly at 60 degrees celsius. Special covers, which fully enclose the mattress, duvet and pillows are available. They prevent the dust mites from getting through.
2. Eliminate or reduce all books and soft toys from the bedroom as they attract the dust mite.
3. Vacuum carpets and soft furnishings twice weekly.
4. Toys: wash furry toys frequently.
5. Ventilate the house well.
6. Flooring: shallow pile carpet or sheer surface floors are preferable e.g. wood or lino.
7. Dust: damp dust all surfaces regularly.

Pets

Dogs and cats are not generally encouraged as pets for children with asthma. Cats tend to produce more symptoms than dogs. If there is already a pet in the home it is best kept outdoors. Pets should never be allowed in the bedroom.




It should be noted that it may take 6 months for cat allergen to be fully eliminated from the house after the cat has been removed.

Viral Infections

Viruses are one of the most common trigger factors for children with asthma. Viruses are common and avoiding them can be very difficult, especially if a child attends crèche/pre-school. Where possible, try to avoid exposure where there are people with known colds/coughs etc.

Fumes

Very high levels of traffic emissions may play an important role in the development of asthma. Tobacco smoke exposure can trigger asthma symptoms and maternal smoking in pregnancy can lead to increase wheeze in infants.





Pollen

Be aware of high pollen counts. There are very effective hay fever treatments available to alleviate the discomfort of symptoms. Discuss these with your doctor or pharmacist if your child has symptoms.

If the pollen count is high:

- Keep windows in the house and car closed
- Your child should wear sunglasses if appropriate
- Your child should wash their hands & face when they come indoors
- Avoid drying clothes outside and/or shake well before bringing inside

Mould

Symptoms of asthma can be triggered by the presence of mould/mildew spores found in damp housing conditions. Avoid condensation in the house. Air the house well every day and avoid drying clothes on radiators.

Food

It is rare for foods to trigger asthma symptoms unless they occur as part of an anaphylactic (allergic) reaction which occurs with foods such as nuts, shellfish, kiwi and egg.

Exercise

Teachers should be informed if your child has asthma. Ensure that there is always a reliever available for use at school. Many teachers will be willing to supervise the administration of medication, however schools vary in their policies on asthma medication. A booklet for teachers on asthma is available from the Asthma Society of Ireland.

1. Children should warm up properly before sports.
2. Wear a scarf around the nose and mouth when exercising outdoors on cold dry days.
3. Swimming is usually a very good form of exercise for asthmatics because the air is warm and moist. However very occasionally chlorine can cause symptoms to worsen.
4. It is useful to give the child their reliever 15 minutes before planned exercise but check with your doctor first.



It is obviously very important that children with asthma lead a normal life and participate in sports and exercise. It is reassuring to know that many famous and successful sports stars have asthma.





Medical treatment of asthma

Always follow your doctor's advice and read the instruction leaflet that comes with the medicine.

The following are the main forms of asthma management

- Relievers: bronchodilators
- Controllers: inhaled steroids
- Combinations: inhaled steroid and a long acting reliever together in one inhaler
- Leukotriene receptor antagonists (LTRA)
- Oral Steroids


Relievers

Relievers work by relaxing the muscles surrounding the airways and allowing them to open up.

They can be:

- Short-acting (act quickly and for a short time)
- Long-acting (work over 12 hours and are used in combination with inhaled steroids)

Short-acting relievers (bronchodilators) are commonly referred to as the 'blue inhaler' and should be taken when your child has asthma symptoms.



If the short-acting reliever inhaler is needed for symptom relief more than 2 times per week, consult with your child's doctor.

Examples of short-acting relievers are Bricanyl®, Salamol® and Ventolin®.

Long-acting relievers are usually taken in the morning and at night. Short-acting relievers are usually taken on an as-needed basis or before exercise.

Another type of reliever is Atrovent® (ipratropium). It is not as fast acting as the other relievers. It is taken up to 4 times per day.


Controllers

Preventers (inhaled steroids) are commonly referred to as the 'brown inhaler'. They reduce the inflammation and swelling in the lining of the airways. They do not provide immediate relief from symptoms, however, they play a very important role in preventing symptoms.





They need to be taken every day in order to be effective, even when your child is feeling well. Some take 2 weeks to become fully effective.



If the preventer is working correctly, your child should not need to use their reliever regularly. Examples of preventers are Beclazone®, Becotide®, Flixotide® and Pulmicort®.

Combinations

These inhalers contain both an inhaled preventer (reducing inflammation) and a long acting reliever (open airways for 12 hours).

These medications are taken regularly and help ensure a simple and effective treatment regime.

Examples are Seretide® and Symbicort®.

Leukotriene receptor antagonists

These are medications that reduce inflammation and spasm in the airways. They are available in tablet form, which can be chewable, or in granules for smaller children.

Oral steroids

In situations where asthma becomes worse, your child may be prescribed a course of steroid tablets. Short courses of oral steroids rarely cause problems. They are very effective at bringing asthma under control and can prevent hospitalisation.

Side effects of asthma medications

Asthma drugs are generally well tolerated and have well documented safety profiles.

Common side effects of the fast-acting **relievers** include a fast heartbeat, hyperactive behaviour and a hand tremor. These effects tend to be felt at the beginning of the treatment or with overuse. The main side effects associated with Atrovent are headaches, dizziness, tummy disturbances, dry mouth, tightening of the chest, cough or some irritation of the airway immediately after taking the inhaler.

Inhaled steroids can sometimes cause hoarseness, sore throat and oral thrush. These side effects may be avoided by mouth rinsing or teeth washing after inhalation. In the case where facemasks are being used, the child's face must also be cleaned.





Parents often worry about inhaled steroids and confuse them with oral steroids. In fact **inhaled steroids** are generally very well tolerated.

A full list of the medicine's side effects can be found in the patient information leaflet that comes with the medicine. Talk to your doctor if you have any concerns.

Complimentary therapy


There is very little evidence to suggest that any form of alternative/complementary treatments help improve asthma symptoms. Eg: Buteyko, reflexology, homeopathy, acupuncture.

It is **very important** to continue to give your child their prescribed medications and to inform your doctor should you be considering this form of treatment.

Inhaler devices and your child

Generally medication in asthma is given through an inhaler. This ensures much smaller doses and that the treatment goes directly to the lungs.

The inhaler device your doctor recommends will depend on your child's age and their ability to use it well. It is important to ask questions and to be confident on how to use your child's inhaler to administer the prescribed medications. Your doctor or practice nurse will demonstrate how to use the device chosen for your child. There is also a DVD available from the Asthma Society of Ireland on 'How to use your inhaler correctly', or visit www.asthmasociety.ie for demonstrations online.



Infants

- Aerosol inhaler with either Aerochamber® or Babyhaler® with mask

Pre-school children (2-4 years)

- Aerosol inhaler with large volume spacer (Aerochamber®, Babyhaler® or Volumatic®).





School age children (4 years +)

- Dry powder inhaler e.g. Diskus[®], Turbohaler[®].
- Aerosol with large volume spacer e.g. Aerochamber[®], Babyhaler[®], or Volumatic[®].
- Breath activated inhaler e.g. Autohaler[®], Easi-Breathe[®].

Tips for giving your child their inhaler

– How to make it easier:

- In the case of small children and infants, make it fun i.e. singing favourite songs, counting to 10 etc.
- Praise and Encouragement
- Patience – give the child a chance to get used to their new inhaler
- Try to make as little fuss as possible. Smile and stay relaxed while your child is taking their inhaler
- Make it part of their routine
- If taking the inhaler through a spacer device, your child may wish to ‘decorate’ their device with their favourite character’s stickers/pictures
- If in doubt your doctor, practice nurse, or pharmacist will be eager to advise

Nebulisers

Nebulisers are machines which deliver a greater dose of the medication to the lungs. Nebulisers are rarely needed in well-controlled asthmatics and are usually seen only in emergency situations or where a child is unable to use a spacer device.

Asthma and Rhinitis

A high percentage of people with asthma also experience symptoms of rhinitis.

Rhinitis is an inflammation of the lining of the nose.

Symptoms may include:

- Sneezing
- Itchy, blocked or runny nose
- Itchy throat, mouth and inner ear
- Headaches
- Loss of concentration
- Feeling generally unwell
- It can feel like a common cold but lasts much longer

Treatments can include antihistamines and nasal preparations – consult your pharmacist or doctor if your child has these symptoms.





Knowing when your child's asthma becomes worse?

As all caregivers know, it is easy to tell when your child is not feeling well.

When your child's asthma is getting worse and you know the signs to look for, symptoms can be quick and simple to manage. Here are some simple tell-tale signs that will help you spot when your child's asthma is getting worse:

- Coughing or wheezing when they get up in the morning or during the night
- Opting out of exercise or not exercising as much as they normally would
- Requesting to use their blue reliever more than normal

If you think your child's asthma control is getting worse, give your doctor or nurse a call.

The future for asthma

Asthma can be successfully managed with regular medications and environmental measures. Fortunately, as many as two thirds of childhood asthma sufferers will grow out of their symptoms. Of the remainder, a significant proportion will improve and require less medications.

Occasionally asthma symptoms resolve in the teenage years only to reoccur in later adulthood. At present we have no accurate way of predicting who will and who will not outgrow their asthma. Asthma remains the subject of constant research and study.

Asthma attack – how can you help?

Asthma attacks can be very frightening so it is important to familiarise yourself with some simple steps to avoid panic should it occur to your child.

Don't be afraid of causing a fuss, even at night. Call a doctor or an ambulance if you have any doubts about your child's condition, or if they are becoming distressed, unable to talk or are getting exhausted. If your child is admitted to hospital or an accident and emergency department because of their asthma, it is important to take their details of treatment with you. You should make an appointment with your doctor or nurse after your child is discharged from hospital, so that you can review his/her asthma treatment to avoid this situation happening again.





For advice on emergency relief from an asthma attack, check out **The Five Minute Rule** on the back of this booklet. The Five Minute Rule has been developed by the Asthma Society of Ireland and a number of Respiratory Consultants in Ireland.

Notes:





Medication Plan

You have been prescribed _____.

Take this _____ in the morning and _____ in the evening.

If you need extra relief take _____ puff of _____

when you need it.

Contact _____ on phone number _____

if you need more advice.

Useful numbers for you

www.asthmasociety.ie

Phone: 1850 44 54 64

Your Doctor or Nurse

Phone: _____

Other booklets available:

- A guide to COPD
- A guide to Asthma



About the author

Dr Peter Greally MD, FRCPI, FRCPCH, DCH graduated from UCD in 1984. He began practice in Paediatrics in 1985 having trained at Dublin, London & Pittsburgh. He has been a member of the Academic Staff in Trinity College Dublin since 1994 and a Consultant Paediatrician also since 1994.



Dr Greally specialises in Children's Respiratory and Allergic Disorders. He works at the National Children's Hospital, Tallaght, Dublin 24, Our Lady's Hospital for Sick Children, Dublin 12 and has a private clinic at Suite 12, Charlemont Clinic, Dublin 2.

He is a Board Member and former Medical Chairperson of Asthma Society of Ireland, a Past Member Board of Governors AMNCH Tallaght, Past Chairman, Division of Paediatrics, AMNCH, Tallaght. He is also Past Secretary of the Irish Paediatric Association.

Dr Greally, is currently a Member of the following: Medical & Scientific Committee Cystic Fibrosis Association of Ireland, Board Member of the Cystic Fibrosis Registry of Ireland, Member of the Irish Thoracic Society (ITS), Founder member of the Paediatric subgroup ITS, Fellow College of Paediatrics and Member of the American Thoracic Society- (Allergy & Immunology Assembly Member).

You can contact Dr Greally by email at dora.daly@amnch.ie or by phone in the Charlemont clinic on 01-414-2188.

The Five Minute Rule

For emergency relief from an asthma attack.

1. Ensure the reliever (blue) inhaler is taken immediately
2. Get your child to sit down and loosen tight clothing
3. Do not put your arm around them
4. Encourage them to breathe slowly and calmly
5. Stay calm and reassure your child

If there is no immediate improvement, continue to take the reliever inhaler every minute for five minutes, or until symptoms improve: two puffs if MDI/evohaler or one puff if turbohaler.

If symptoms do not improve in five minutes, or if you are in doubt, call 999 or a doctor urgently.

- Continue to give reliever inhaler until help arrives or symptoms improve
- Take your child's treatment details to the hospital or accident and emergency department

Don't be afraid of causing a fuss, even at night. Call the doctor or an ambulance if you have any doubts about your child's condition or if they are becoming distressed, unable to talk or are getting exhausted. If your child is admitted to hospital or an accident and emergency department because of their asthma, it is important to take details of their treatment with you. You should also make an appointment with your doctor or nurse after your child is discharged from hospital, so that you can review his/her asthma treatment to avoid this situation happening again.

The Five Minute Rule has been developed by the Asthma Society of Ireland and a number of Respiratory Consultants in Ireland.