

Boehringer Ingelheim is pleased to provide you with this Educational Pack, which has been developed to give practical and relevant information on the appropriate use of Pradaxa®. The pack includes:

- Pradaxa® 150mg – Summary of Product Characteristics
- Pradaxa® 110mg – Summary of Product Characteristics
- Prescriber Guide – this addresses recommendations for the use of Pradaxa® in order to minimise the risk of bleeding
- Patient Alert Card

To order additional copies of the Patient Alert Card please go to: www.pradaxa.co.uk/DVT-PEeducationalpack

You can also order or download this Educational Pack.

References:

1. Boehringer Ingelheim. Pradaxa® 150mg hard capsules Summary of Product Characteristics.
2. Boehringer Ingelheim. Pradaxa® 110mg hard capsules Summary of Product Characteristics.

Prescribing Information (DVT/PE – UK) PRADAXA® (dabigatran etexilate)

Capsules containing 110 mg or 150 mg dabigatran etexilate (as mesilate) **Action:** Direct thrombin inhibitor. **Indication:** Treatment of deep vein thrombosis (DVT) and pulmonary embolism (PE), and prevention of recurrent DVT and PE in adults (DVT/PE). **Dose and Administration:** Renal function should be assessed by calculating CrCL prior to initiation to exclude patients with severe renal impairment (CrCL < 30 mL/min). Recommended daily dose 300 mg taken as one 150 mg capsule twice daily following treatment with parenteral anticoagulant for at least 5 days. Duration of treatment should be individualised after careful assessment of the treatment benefit against risk for bleeding. Short duration of therapy (at least three months) should be based on transient risk factors (e.g. recent surgery, trauma, immobilisation) and longer durations should be based on permanent risk factors or idiopathic DVT or PE. In case of intolerance to dabigatran, patients should be instructed to immediately consult their doctor. For patients aged 80 years or above, or those receiving concomitant verapamil, the recommended daily dose is Pradaxa 220 mg taken as 110 mg twice daily. Pradaxa and verapamil should be taken at the same time. For the following patient groups, the daily dose of 300 mg or 220 mg should be selected based on an individual assessment of the thromboembolic risk and risk of bleeding: aged 75 – 80 years; with moderate renal impairment (CrCL 30-50 mL/min); with gastritis, oesophagitis or gastroesophageal reflux; other risk of increased bleeding. Close clinical surveillance is recommended in patients with renal impairment. Use is contraindicated in patients with severe renal impairment (CrCL < 30 mL/min). In all patients assess renal function by calculating CrCL prior to initiation to exclude patients with severe renal impairment. Renal function should also be assessed when a decline in renal function is suspected. Additionally in patients > 75 years or with mild to moderate renal impairment, renal function should also be assessed at least once a year or more frequently as needed in certain clinical situations when it is suspected that the renal function could decline or deteriorate. Patients with an increased risk of bleeding: closely monitor clinically looking for signs of bleeding or anaemia. A coagulation test may help identify increased risk patients. No dose adjustment required but close clinical surveillance in patients < 50 kg. Not recommended if liver enzymes > 2 Upper Limit of Normal (ULN). If switching from Pradaxa to parenteral anticoagulant wait 12 hours after the last dose of Pradaxa; if switching from parenteral anticoagulant to Pradaxa discontinue the parenteral anticoagulant and start Pradaxa 0-2 hours prior to the time that the next dose of the alternate therapy would be due, or at the time of discontinuation in case of continuous treatment; if switching from Pradaxa to VKA adjust the starting time of the VKA based on CrCL; if switching from VKA to Pradaxa stop VKA and give Pradaxa once INR <2.0. Cardioversion patients can stay on Pradaxa whilst being cardioverted. The safety and efficacy of Pradaxa in ages less than 18 years have not been established in the indication. Pradaxa can be taken with or without food. Pradaxa should be swallowed as a whole with a glass of water to facilitate delivery to the stomach. Patients should be instructed not to open the capsule as this may increase the risk of bleeding. **Contraindications:** Hypersensitivity to any component; severe renal impairment (CrCL < 30 mL/min); active clinically significant bleeding; lesion or condition, if considered a significant risk factor for major bleeding. This may include current or recent gastrointestinal ulceration, presence of malignant neoplasms at high risk of bleeding, recent brain or spinal injury, recent brain, spinal or ophthalmic surgery, recent intracranial haemorrhage, known or suspected oesophageal varices, arteriovenous malformations, vascular aneurysms or major intraspinal or intracerebral vascular abnormalities; concomitant treatment with any other anticoagulants e.g. unfractionated heparin (UFH), low molecular weight heparins (enoxaparin, dalteparin etc), heparin derivatives (fondaparinux etc), oral anticoagulants (warfarin, rivaroxaban, apixaban etc) except under specific circumstances of switching anticoagulant therapy or when UFH is given at doses necessary to maintain an open central venous or arterial catheter; hepatic impairment or liver disease expected to have any impact on survival; concomitant systemic ketoconazole, cyclosporine, itraconazole, dronedarone; prosthetic heart valves requiring anticoagulant treatment. **Warnings and Precautions:** Not recommended if liver enzymes > 2 ULN. Haemorrhagic risk: Close clinical surveillance (signs of bleeding or anaemia) is recommended throughout the treatment period, especially when haemorrhagic risk is increased or risk factors combined. Factors which may increase haemorrhagic risk: age ≥ 75 years; moderate renal impairment (CrCL 30 – 50 mL/min); P-glycoprotein inhibitor co-medication; body weight <

50 kg; acetylsalicylic acid (aspirin); NSAID; clopidogrel; selective serotonin re-uptake inhibitors (SSRIs) or selective serotonin norepinephrine re-uptake inhibitors (SNRIs); other drugs which may impair haemostasis; diseases/procedures associated with a risk of bleeding such as coagulation disorders, thrombocytopenia or functional platelet defects, recent biopsy, major trauma, bacterial endocarditis, oesophagitis, gastritis or gastroesophageal reflux. Concomitant use of ticagrelor. The measurement of dabigatran related anticoagulation may be helpful to avoid excessive high exposure to dabigatran in the presence of additional risk factors. Patients who develop acute renal failure must discontinue Pradaxa. If severe bleeding occurs, discontinue treatment and investigate the source of the bleeding. Avoid or use with caution medicinal products which may increase the risk of haemorrhage. The use of fibrinolytic medicinal products for the treatment of acute ischaemic stroke may be considered if the patient presents with a dTT, ECT or aPTT not exceeding the ULN according to the local reference range. Avoid concomitant administration with P-gp inducers. Patients on dabigatran etexilate who undergo surgery or invasive procedures are at increased risk for bleeding therefore surgical interventions may require the temporary discontinuation of dabigatran etexilate; prescribers should consult the Summary of Product Characteristics for further information. Procedures such as spinal anaesthesia may require complete haemostatic function. The risk of spinal or epidural haematoma may be increased in cases of traumatic or repeated puncture and by the prolonged use of epidural catheters. After removal of a catheter, an interval of at least 2 hours should elapse before the administration of the first dose of dabigatran etexilate; these patients require frequent observation for neurological signs and symptoms of spinal or epidural haematoma. Treat with caution patients at high surgical mortality risk and with intrinsic risk factors for thromboembolic events. Myocardial infarction. Efficacy and safety have not been established for DVT/PE patients with active cancer. Contains Sunset Yellow (E110) which may cause allergic reactions. **Interactions:** Anticoagulants and antiplatelet aggregation medicinal products; P-gp inhibitors e.g. amiodarone, quinidine, verapamil, clarithromycin, ticagrelor co-administration (close clinical surveillance); verapamil co-administration - reduce Pradaxa dose to 220 mg (see above) close clinical surveillance is recommended; caution when co-administered with posaconazole; not recommended for concomitant treatment tacrolimus, protease inhibitors including ritonavir and its combinations with other protease inhibitors; avoid with P-gp inducers e.g. rifampicin, St John's wort, carbamazepine, phenytoin; SSRIs or SNRIs. Dabigatran etexilate and dabigatran are not metabolised by cytochrome CYP450 system, therefore related medicinal product interactions not expected. Pantoprazole and other proton-pump inhibitors (PPI) were co-administered with Pradaxa in clinical trials and concomitant PPI treatment did not appear to reduce the efficacy of Pradaxa. Ranitidine administration together with Pradaxa had no clinically relevant effect on the extent of absorption of dabigatran. **Fertility, pregnancy and lactation:** Avoid pregnancy during treatment. Do not use in pregnancy unless clearly necessary. Discontinue breast-feeding during treatment. **Undesirable effects:** Most commonly reported adverse reactions are bleedings occurring in total in approximately 14.4 % in patients treated for DVT/PE. Bleedings occurred in 19.4% of patients in DVT/PE prevention trial RE-MEDY and in 10.5% of patients in DVT/PE prevention trial RE-SONATE. Common (≥ 1/100 to <1/10): epistaxis; gastrointestinal haemorrhage; dyspepsia; rectal haemorrhage; skin haemorrhage; genitourological haemorrhage, including haematuria. Prescribers should consult the Summary of Product Characteristics for further information on side effects. **Pack sizes and NHS price:** 110 mg 60 capsules £65.90 150 mg 60 capsules £65.90 **Legal category** POM **MA numbers:** 110 mg EU/1/08/442/007 (60 capsules) 150 mg EU/1/08/442/011 (60 capsules) **Marketing Authorisation Holder:** Boehringer Ingelheim International GmbH, Binger Str. 173, D-55216 Ingelheim am Rhein, Germany. Prescribers should consult the Summary of Product Characteristics for full prescribing information. **Prepared in** June 2014.

Adverse events should be reported. Reporting forms and information can be found at www.mhra.gov.uk/yellowcard. Adverse events should also be reported to Boehringer Ingelheim Drug Safety on 0800 328 1627 (trefophone).



Date of preparation: June 2014 Job code: UK/DBG-141127



PRADAXA® (DABIGATRAN ETEXILATE) EDUCATIONAL PACK

For treatment of deep vein thrombosis (DVT) and pulmonary embolism (PE), and prevention of recurrent DVT and PE in adults.^{1,2}