

Please make sure that you select the correct MabThera® (rituximab) formulation and strength

SUBCUTANEOUS INJECTION

**MabThera 1400 mg
ONLY FOR USE IN NHL***



Pink flip-off cap

Withdraw directly from vial and administer by subcutaneous injection



Check for the specific MabThera SC packaging characteristics before use:

Red labelling: **'Only for subcutaneous use', 'solution for subcutaneous injection'** and **'subcutaneous'**

MabThera SC is not indicated as once-weekly monotherapy in patients with relapsed/refractory follicular lymphoma

*NHL = non-Hodgkin's lymphoma.

Please refer to the MabThera Summary of Product Characteristics for further information: www.medicines.org.uk or www.medicinesauthority.gov.mt See overleaf for how to use the peel-off part of the label

INTRAVENOUS INJECTION

**MabThera 100 mg concentrate for solution for infusion
MabThera 500 mg concentrate for solution for infusion
For use in all MabThera-approved indications***




Dilute with 0.9% NaCl or 5% glucose and administer by intravenous infusion



Use the peel-off part of the label to ensure the correct formulation and strength is administered to your patient, as illustrated below

**MabThera 1400 mg solution for subcutaneous injection
ONLY FOR USE IN NHL***



1. The MabThera 1400 mg vial label has a removable part

2. Remove the peel-off part

3. Stick the peel-off part on to the syringe

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Reporting forms and information can be found at www.medicinesauthority.gov.mt/adrportal. Adverse events should also be reported to Roche Products Limited. Please contact Roche Drug Safety Centre by emailing welwyn.uk_dsc@roche.com or calling +44(0)1707 367554. As MabThera is a biological medicine, healthcare professionals should report adverse reactions by brand name and batch number.