



This card contains important safety information that you need to be aware of before you are given Enbrel and during treatment with Enbrel. If you do not understand this information, please ask your doctor to explain it to you.

- Show this card to any doctor involved in your treatment.
- See the Enbrel package leaflet for more information.
- Keep this card with you for 2 months after the last Enbrel dose, since side effects may occur after your last dose of Enbrel.

Infections

Enbrel may increase your risk of getting infections, which could be serious.

- You should not use Enbrel if you have an infection. If you are not sure, ask your doctor.
- If you develop symptoms suggestive of infections, such as fever, persistent cough, weight loss, or listlessness, seek medical attention immediately.
- You should be evaluated for tuberculosis (TB). Ask your doctor to record the dates and the results of your last screening for TB below:

Test: Test:

Date: Date:

Results: Results:

- Please ask your physician to list your other medications that may increase your risk of infection.

.....

.....

.....

Congestive Heart Failure

- If you develop symptoms suggestive of congestive heart failure or worsening of existing congestive heart failure, such as shortness of breath, swelling of ankles, persistent cough or fatigue, seek medical attention immediately.

Other Information *(please complete)*

Patient's Name:

Doctor's Name:

Doctor's Phone:



PATIENT ALERT CARD

This card contains important safety information that you need to be aware of before you are given Enbrel and during treatment with Enbrel. If you do not understand this information, please ask your doctor to explain it to you.

- Show this card to any doctor involved in your treatment.
- See the Enbrel package leaflet for more information.
- Keep this card with you for 2 months after the last Enbrel dose, since side effects may occur after your last dose of Enbrel.

Infections

Enbrel may increase your risk of getting infections, which could be serious.

- You should not use Enbrel if you have an infection. If you are not sure, ask your doctor.
- If you develop symptoms suggestive of infections, such as fever, persistent cough, weight loss, or listlessness, seek medical attention immediately.
- You should be evaluated for tuberculosis (TB). Ask your doctor to record the dates and the results of your last screening for TB below:

Test: Test:

Date: Date:

Results: Results:

- Please ask your physician to list your other medications that may increase your risk of infection.

.....

.....

.....

Congestive Heart Failure

- If you develop symptoms suggestive of congestive heart failure or worsening of existing congestive heart failure, such as shortness of breath, swelling of ankles, persistent cough or fatigue, seek medical attention immediately.

Other Information *(please complete)*

Patient's Name:

Doctor's Name:

Doctor's Phone: