m o

Gather your supplies

Find a clean, well-lit surface and gather:




Blue needle cap

A new, pre-filled Benepali®pen autoinjector (remove from carton stored in refrigerator)

A single-use pre-filled pen contains one 50 mg dose of Benepali.

Sharps disposal

container



Inspect carefully

Inspect the Benepali® pen

Do not use the pen past the expiration date.

Do not use if dropped on a hard surface as components inside may be broken.

Do not shake the pen.

Do not use if the blue needle cap is not securely attached.

Inspect the medication

Look through the viewing window. The medication should appear clear or slightly opalescent, colourless or pale yellow, and may contain small white or almost transparent particles of protein.

Do not use the medication if it is discoloured, cloudy, or if particles other than those described above are present.

Let medication reach room temperature

Allow 30 minutes

Leave the Benepali® pen out at room temperature for at least 30 minutes. This makes the medication easier and more comfortable to inject.

Keep out of reach of any children.

Do not use heat sources such as a microwave or hot water to warm the medication

Do not remove the blue needle cap.



Wait 30 minutes before using medication

Wash hands Use soap and water.



Clean injection site

Wipe the skin at the injection site with an alcohol swab.

Do not touch this area again before injecting.



Remove needle cap

Pull the blue needle cap straight off.

Do not twist or bend the blue needle cap while removing which may damage the needle.

Never recap the needle.

Dispose of cap in the sharps



Stretch skin, position pen

Gently stretch the skin at the injection site.

Position the Benepali® pen at approximately 90 degrees to the skin.

Do not pinch the skin.

Listen for 2 clicks while pressing



Press firmly to start

Firmly grip the Benepali® pen and press down into the site to start the injection (there is no button to push).

The pen will **first click** when the injection begins - do not lift the pen

Continue to hold the pen firmly pressed into the site.

Listen for a second click.

Keep pressing for 15s



Once you hear the second click,

to press into the injection site to

Do not release pressure against the

injection site or move the Benepali®

pen before injection is complete.

complete proper injection.

Check yellow



Count to 15 after 2nd click Remove and confirm

Lift empty pen off skin. count slowly to 15 while continuing

The needle guard will completely cover the needle.

> **Check** that the yellow plunger rod extends through the viewing window to confirm that the full dose has been delivered.

Choose where to inject

Select your injection site

The Benepali® pen is for injection under the skin (subcutaneous injection). It should be injected into the thigh, abdomen (at least 5 cm away from the belly button), or back of the upper arm.

Use a different site for each injection. Each new injection site should be given at least 3 cm (or the width of 3 fingers) from an old site.

Do not inject into areas that are red, hard, bruised or tender, or into scars or stretch marks.

If you have psoriasis, do not inject into any raised, thick, red or scaly skin patches (psoriasis skin lesions)

Dispose of pen properly



Do not throw the sharps container in household bin or recycle it.

dispose of a filled sharps container.

Always keep the container out of the sight and reach of children

Check injection site

If there is bleeding, press a cotton ball or gauze pad over the injection site.

If needed, cover with a plaster.

Do not rub the injection site.



Questions?

efer to your *Instructions for Use* and package

eporting of Side Effects

possible side effects not listed in the package leaflet. You can also report side effects directly in the package leaflet. ormation on the safety of this medicine.

This medicine is subject to dditional monitoring. Reporting suspected adverse events or reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse events or reactions

Adverse events can be reported directly via:

Malta

Pharma.MT limited Tel: +356 213 37008/9

Malta Medicines Authority ADR Reporting details:

www.medicinesauthority.gov.mt/ adrportal

After the injection

Read the *Instructions For Use* that come with Benepali* completely before using the Benepali* pen and each time you get a prescription refill as there may be new information. This *Quick Reference Guide* is not intended to replace those instructions. **Do not** try to give yourself the injection unless you have received training from your doctor or nurse.

(PFPen) EN v 1.1. Approved: BEN-PAN-0018a 02/2016