

## What should I know about Xarelto?

- Xarelto thins the blood, which prevents from getting dangerous blood clots
- Xarelto must be taken exactly as prescribed by the doctor. To ensure optimal protection from blood clots, **never skip a dose**
- You must not stop giving/taking Xarelto without first talking to the doctor, as the risk of blood clots may increase
- Tell the healthcare provider about any other medicines the child is/you are currently taking, took recently or intend to start taking, before starting Xarelto
- Tell the healthcare provider that the child is/you are taking Xarelto before any surgery or invasive procedure

## When should I seek advice from a healthcare provider?

When taking a blood thinner such as Xarelto, it is important to be aware of its possible side effects. Bleeding is the most common side effect. Do not start giving Xarelto to the child/taking Xarelto if you know the child is/you are at risk of bleeding, without first discussing this with the doctor.

Tell the healthcare provider straight away if the child has/you have any signs or symptoms of bleeding, such as the following:

- Pain
- Swelling or discomfort
- Headache, dizziness or weakness
- Unusual bruising, nosebleeds, bleeding of gums, cuts that take a long time to stop bleeding
- Menstrual flow or vaginal bleeding that is heavier than normal

- Blood in the urine which may be pink or brown; red or black stools
- Coughing up blood, or vomiting blood or material that looks like coffee grounds

## How do I give Xarelto to the child?/How do I take Xarelto?

To ensure optimal protection, Xarelto 1 mg/mL must be given/taken with feeding (breast milk or formula feeding) or with a meal.

It can be administered through a nasogastric or gastric feeding tube.



# Patient Alert Card

Xarelto® (rivaroxaban)  
1 mg/mL granules for oral suspension

### Information to caregivers/patients:

- **Keep this card with you at all times**
- **Present this card to every physician or dentist prior to treatment of the patient**

Please watch the educational video showing how to prepare and administer the oral suspension.



[go.bayer.com/junior](https://go.bayer.com/junior)

## The patient who is under anticoagulation treatment with Xarelto (rivaroxaban)

---

\_\_\_\_\_

Name

\_\_\_\_\_

Other medicines/conditions

\_\_\_\_\_

Address

\_\_\_\_\_

Birth date

\_\_\_\_\_

Weight

## In case of emergency, please notify:

---

\_\_\_\_\_

Doctor's name

\_\_\_\_\_

Doctor's phone

\_\_\_\_\_

Doctor's stamp

## Please also notify:

---

\_\_\_\_\_

Name

\_\_\_\_\_

Phone

\_\_\_\_\_

Relationship

## Information for healthcare providers:

---

- INR values should not be used as they are not a dependable measure of the anticoagulant activity of Xarelto