

Keep this information with you and share it with other healthcare professionals involved in your medical care or treatment.

Your Name:

Doctor's Name (who prescribed Olumiant):

Doctor's phone number:

Patient Alert Card

Information for Patients about OLUMIANT® (baricitinib)

This document contains important information you should be aware of before and during treatment with Olumiant.

Pregnancy

- Do not take Olumiant if you are pregnant or suspect you may be pregnant
- Use effective contraception while taking Olumiant (and for 1 week after, if you stop treatment)
- Tell your doctor immediately if you become (or wish to become) pregnant

Infections

Olumiant may make an existing infection worse or increase the chance of you getting a new infection or increase the chance of viral reactivation. Inform your doctor immediately if you get symptoms of infection, such as:

- Fever, wounds, feeling more tired than usual, or dental problems
- A cough that won't go away, night sweats, and weight loss. These could be symptoms of tuberculosis (an infectious disease of the lungs)
- A painful skin rash with blisters. This could be a sign of a herpes zoster infection

Blood fat

Your doctor may check for levels of fat in the blood, such as cholesterol, while you are taking Olumiant.

Blood clots

Olumiant may cause a condition in which a blood clot forms in your leg that may travel to your lungs. Inform your doctor immediately if you experience any of the following symptoms:

- Swelling or pain in one leg
- Warmth or redness in one leg
- Shortness of breath which is unexpected
- Rapid breathing
- Chest pain