

## EMA confirms measures to minimise risk of serious side effects with Janus kinase inhibitors for chronic inflammatory disorders

14.11.2022 | Circular Number P14/2022

### Information on Janus kinase inhibitors

- The Janus kinase (JAK) inhibitors subject to this review<sup>1</sup> are used to treat several chronic inflammatory disorders (rheumatoid arthritis, psoriatic arthritis, juvenile idiopathic arthritis, ankylosing spondylitis, ulcerative colitis, atopic dermatitis, and alopecia areata).
- The active substances in these medicines work by blocking the action of enzymes known as Janus kinases. These enzymes play an important role in the process of inflammation that occurs in these disorders. By blocking the enzymes' action, the medicines help reduce the inflammation and other symptoms of these disorders.

Some JAK inhibitors (Jakavi and Inrebic) are used to treat myeloproliferative disorders; the review did not include these medicines. The review also did not cover the use of Olumiant in the short-term treatment of COVID-19, which is under assessment by EMA.

The following products are authorised via centralised procedure.

Active Ingredients	Product Name	Pharmaceutical Form	Classification	Authorisation Number	MAH/license holder
Baricitinib	Olumiant	Film-coated tablet	POM	EMEA/H/C/004085	Eli Lilly Nederland B.V.
Tofacitinib	Xeljanz	Film-coated tablet	POM	EMEA/H/C/004214	Pfizer Europe MA EEIG

### Information from the EMA about the safety concern

- The review of JAK inhibitors in the treatment of inflammatory disorders was initiated at the request of the European Commission (EC) under [Article 20 of Regulation \(EC\) No 726/2004](#).
- The review was carried out by the Pharmacovigilance Risk Assessment Committee (PRAC), the Committee responsible for the evaluation of safety issues for human medicines, which made a set of recommendations. The PRAC recommendations were forwarded to the EMA's human medicines committee (Committee for Medicinal Products for Human Use (CHMP)), responsible for questions concerning medicines for human use,

<sup>1</sup>Olumiant (baricitinib), and Xeljanz (tofacitinib).

which adopted the Agency's opinion. CHMP has endorsed the measures recommended by the PRAC to minimise the risk of serious side effects with JAK inhibitors.

- The recommendations follow a review of available data, including the final results from a clinical trial of the JAK inhibitor Xeljanz (tofacitinib) and preliminary findings from an observational study involving Olumiant. The review also included advice from an expert group of rheumatologists, dermatologists, gastroenterologists and patient representatives.
- These medicines should be used in the following patients only if no suitable treatment alternatives are available: those aged 65 years or above, those at increased risk of major cardiovascular problems (such as heart attack or stroke), those who smoke or have done so for a long time in the past and those at increased risk of cancer.
- JAK inhibitors should be used with caution in patients with risk factors for blood clots in the lungs and in deep veins (venous thromboembolism, VTE) other than those listed above. Further, the doses should be reduced in patient groups who are at risk of VTE, cancer or major cardiovascular problems, where possible.

### **Information to Healthcare Professionals**

- An EMA review has found that, compared with TNF-alpha inhibitors, JAK inhibitors used to treat chronic inflammatory disorders (rheumatoid arthritis, psoriatic arthritis, juvenile idiopathic arthritis, axial spondyloarthritis, ulcerative colitis, atopic dermatitis and alopecia areata) are linked to a higher risk of major adverse cardiovascular events (MACE), venous thromboembolism (VTE), malignancy, serious infections and all-cause mortality.
- The review confirmed Xeljanz increases the risk of major cardiovascular problems, cancer, VTE, serious infections and death due to any cause when compared with medicines belonging to the class of TNF-alpha inhibitors. EMA has now concluded that these safety findings apply to all approved uses of JAK inhibitors in chronic inflammatory disorders (rheumatoid arthritis, psoriatic arthritis, juvenile idiopathic arthritis, axial spondyloarthritis, ulcerative colitis, atopic dermatitis and alopecia areata). The review included the final results from an open-label clinical trial (ORAL Surveillance study)<sup>2</sup> of the JAK inhibitor Xeljanz (tofacitinib) in patients with rheumatoid arthritis and cardiovascular risk factors which found a higher risk of these events with Xeljanz than with TNF-alpha inhibitors.
- Preliminary findings from an observational study (B023) involving another JAK inhibitor, Olumiant (baricitinib), also suggest an increased risk of MACE and VTE in patients with rheumatoid arthritis treated with Olumiant compared with those treated with TNF-alpha inhibitors.
- EMA concluded that the identified risks apply to all JAK inhibitors<sup>3</sup> approved for the treatment of chronic inflammatory disorders. These medicines should only be used in the

---

<sup>2</sup> Ytterberg SR, et al. Cardiovascular and cancer risk with tofacitinib in rheumatoid arthritis. *New Engl J Med* 2022;386(4):316-326. doi: [10.1056/NEJMoa2109927](https://doi.org/10.1056/NEJMoa2109927)

<sup>3</sup> Olumiant (baricitinib), and Xeljanz (tofacitinib)

following patients if no suitable treatment alternatives are available: those aged 65 years or above, those who are current or past long-time smokers, those with a history of atherosclerotic cardiovascular disease or other cardiovascular risk factors, or those with other malignancy risk factors. Cautious use is also recommended in patients with known risk factors for VTE other than those listed above.

- If JAK inhibitors<sup>4</sup> are needed in patients with these risk factors, a lower dose may be recommended, depending on the medicine, the indication and the specific risk factor. Healthcare professionals should discuss the risks associated with JAK inhibitors with their patients. It is recommended that healthcare professionals carry out periodic examinations of their patients' skin to check for skin cancer, particularly for patients at risk for skin cancer.
- A letter will be sent to all healthcare professionals expected to prescribe these medicines to inform them of the outcome of the review. Full treatment recommendations will be included in the updated summary of product characteristics and the educational material for the respective products.

### **Information to patients**

- JAK inhibitors used to treat chronic inflammatory disorders have been found to increase the risk of major cardiovascular problems (such as heart attack or stroke), cancer, blood clots in the lungs and in deep veins, serious infections and death when compared with TNF alpha inhibitors.
- These JAK inhibitors are used to treat one or more of the following chronic inflammatory disorders: rheumatoid arthritis, psoriatic arthritis, juvenile idiopathic arthritis, axial spondyloarthritis, ulcerative colitis, atopic dermatitis and alopecia areata.
- If you are aged 65 years or above, have an increased risk of major cardiovascular problems or cancer or if you smoke or have done so for a long time in the past, you should only be prescribed these medicines if there are no suitable treatment alternatives for you. If you have certain risk factors, your doctor may reduce the dose of your JAK inhibitor or switch treatment depending on your inflammatory disorder and the JAK inhibitor you are taking to treat it.
- If, at any stage during your treatment, you experience chest pain or tightness (which may spread to arms, jaw, neck and back), shortness of breath, cold sweat, light headedness, sudden dizziness, weakness in arms and legs or slurred speech, contact your doctor immediately. Examine your skin periodically and let your doctor know if you notice any new growths on the skin. If you have any questions about your treatment, speak to your doctor.

For more information please see the European Medicines Agency's [press release](#).

---

## **Reporting Adverse Drug Reactions**

Healthcare professionals and patients are encouraged to maintain vigilance with Janus kinase inhibitors<sup>5</sup>. Suspected Adverse Drug Reactions (side effects) may be reported using the Medicines Authority Form and sending it to: Sir Temi Żammit Buildings, Malta Life Sciences Park, San Ġwann SĠN 3000 **or** online to <http://www.medicinesauthority.gov.mt/adrportal> or to the marketing authorisation holder or their local representatives.

## **Post-Licensing Directorate**

### **Medicines Authority**

*Healthcare professionals and patients are encouraged to regularly check the Medicines Authority website for product safety updates as these are issued on an ongoing basis.*

---

<sup>5</sup>Olumiant (baricitinib), and Xeljanz (tofacitinib).

**Feedback Form**

The Medicines Authority thanks you for the time taken to read this safety circular. The dissemination of safety circulars is an important process whereby Regulatory Authorities can communicate important issues with respect to the safety of medicines, in order to protect and enhance public health.

The Medicines Authority kindly invites your anonymous feedback about the regulatory action being communicated. This may be returned by folding this form (address side up), stapling the ends and then posting (no stamp required).

**Feedback:**

---

---

---

---

---

---

---

---

---

---

We thank you for your interest and look forward to hearing your opinion.

Postage will be paid  
by the Licensee

No postage stamp  
necessary if posted  
in Malta and Gozo

**BUSINESS REPLY SERVICE**

Licence no. 656

Pharmacovigilance Section  
Post-Licensing Directorate  
Medicines Authority  
Sir Temi Żammit Buildings  
Malta Life Sciences Park  
San Ġwann SĠN 3000