

Considerations when ending treatment with Amfexa[®] 5mg, 10 mg and 20 mg Tablets

To prevent unsupervised termination of medication, the procedures for a trial period off medicine should be discussed routinely with patients. Regular assessment and monitoring of behaviour through formal ratings appears to decrease the rate of unsupervised termination of medications.

After years of medication, it is reasonable to offer children and adolescents who have had stable improvement in ADHD symptoms and target behaviours a trial off of medication to determine whether medication is still necessary. Such a trial should be undertaken with close monitoring of core symptoms and function at home, school, and in the community.

Careful supervision is required during dexamfetamine withdrawal since this may unmask depression as well as chronic hyperactivity.

Some patients may require long-term follow-up.

If you have any questions relating to this medication, please contact us at Medice, UK

Telephone: 0204 582 2845

Email: medicalinformation@medice.co.uk

If you need to report any Adverse Drug Reaction please inform the regulatory authority <https://medicinesauthority.gov.mt/reportingadversereactions> or the local representative:

EJ Busuttil Ltd

Busuttil Buildings, Triq I-Ghadam, Central Business District Zone 1

Birkirkara

Tel 00356 21447184

rp@ejbusuttil.com

If you need to report any Adverse Drug Reaction please inform the regulatory authority <https://medicinesauthority.gov.mt/reportingadversereactions> or the local representative:

EJ Busuttil Ltd

Busuttil Buildings, Triq I-Ghadam, Central Business District Zone 1

Birkirkara

Tel 00356 21447184

rp@ejbusuttil.com