

Otsuka **Jinarc®** ▼ (Tolvaptan) Patient Alert Card

Patient's Name:

Date Jinarc® first prescribed:

Doctor's Name:

Hospital name and telephone number:

Important safety information for patients

Jinarc® can affect how your liver works.

Consult your doctor if you experience symptoms of tiredness, loss of appetite, upper abdominal pain or discomfort, fever, dark urine or yellowing of skin or eyes, joint and muscle pain with fever (flu-like syndrome), itching, nausea and vomiting.

Jinarc® can cause severe dehydration.

Drink plenty of fluids to avoid dehydration or excessive water loss and consult your doctor if you are not able to drink fluids by mouth.

Important safety information for emergency carers

Jinarc® may cause liver injury. Blood tests for liver function must be performed periodically (monthly for the first 18 months, then every 3 months of continuing treatment). Therapy should be stopped or discontinued if significant increase of liver enzymes and/or clinical symptoms of hepatic injury persist.

Jinarc® can produce significant increase in urination which may lead to severe dehydration or excessive water loss. Symptoms of dehydration may include increased thirst, dry mouth, feeling tired or sleepy, decreased urination, headache, dry skin, dizziness, rapid heart rate, confusion and poor skin elasticity.

For more info please contact the representative office of the holder license Swixx Biopharma S.M.S.A, tel: +30 214 444 9670, email: medinfo.malta@swixxbiopharma.com