



Vosoritide (VOXZOGO[®]) treatment placemat

Use this placemat on a clean, dry surface, and prepare for your child's vosoritide injection by setting out all of the items indicated. To properly administer vosoritide, follow the instructions in the caregiver dosing and administration guide and patient injection video. **Make sure to have your sharps disposal or puncture-resistant container nearby. Check with your child's doctor, a pharmacist or nurse about the correct way to dispose of sharps.**

Alcohol Pads




Solvent Syringe



Vosoritide Vial


Cap colours vary with different doses



Side view

Remove cap and twist solvent needle on here


Solvent Needle



Twist this end onto solvent syringe

Remove cap before inserting needle into vosoritide vial


Injection Syringe



Needle cap

Plunger cap

Adhesive Bandage or Gauze



Keep these and all medications and medical equipment out of the reach of children.



Scan with your phone or visit vosoritideinjection.com to watch video about how to inject vosoritide.

This material is only for caregivers of patients who have been prescribed vosoritide. It is an additional information resource that should be read alongside the package leaflet. Always refer to the package leaflet before beginning treatment with vosoritide. If you have any questions, please discuss them with your child's doctor.

Reporting of side effects: If your child gets any side effects, talk to your child's doctor, or nurse. This includes any possible side effects not listed in the package leaflet. By reporting side effects you can help provide more information on the safety of this medicine.

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects directly to BioMarin via drugsafety@bmrn.com; Fax: +1-415-532-3144 or Tel: +1-415-506-6179. Reporting of side effects is also possible via national reporting systems.

Please see the important safety information on the reverse, and the accompanying package leaflet.

Suspected Adverse Drug Reactions (side effects) or medication errors may be reported using the Medicines Authority ADR reporting form, which is available online at <http://www.medicinesauthority.gov.mt/adrportal>, and sent by post or email to; P: Pharmacovigilance Section at Post-Licensing Directorate, Medicines Authority, Sir Temi Zammit Buildings, Malta Life Sciences Park, San Ġwann SĠN 3000 E: postlicensing.medicinesauthority@gov.mt

What is vosoritide?

Vosoritide is a prescription medicine used for the treatment of achondroplasia in children 4 months of age and older whose bones are still growing.

What do you need to know before you use vosoritide?

Do not use vosoritide

- If your child is allergic to vosoritide or any of the other ingredients of this medicine listed in the package leaflet

Warnings and precautions

Talk to your child's doctor before using vosoritide:

- If your child has significant heart disease or blood pressure problems
- If your child is taking or has recently taken medicines that lower blood pressure

If any of these apply to your child or you are unsure, talk to your child's doctor before using vosoritide.

Vosoritide can lower blood pressure. As a result, your child may feel dizzy, nauseous, or tired. Blood pressure usually returns to normal within 90 minutes of vosoritide injection. If these effects occur and are severe, tell your child's doctor. Drinking plenty of fluids at the time of injection may reduce the likelihood of these effects. It is recommended patients eat a light snack and drink enough fluid (e.g. water, milk or juice) about 30 minutes before injection.

Other medicines and vosoritide

Tell your child's doctor if your child is taking, has recently taken or might take any other medicines.

Pregnancy and breast-feeding

If your child is being treated with this medicine and is pregnant or breast-feeding, think they may be pregnant or are planning to have a baby, ask their doctor for advice before using this medicine. The use of this medicine is not recommended during pregnancy and breast-feeding.

Driving, cycling and using machines

This medicine may cause your child to feel dizzy, light-headed, tired or your child may feel sick shortly after their injection. If this happens, they should not drive, ride a bicycle, do physical activities or use machines for around an hour after injection or until they feel better.

Vosoritide contains sodium

This medicine contains less than 1 mmol sodium (23mg) per dose, that is to say essentially 'sodium-free'.

BOMARIN®

Developed and Funded by BioMarin International Ltd.
© 2024 BioMarin International Ltd. All rights reserved.
EUCAN-VOX-00237 April 2024

Important notes

If you use more vosoritide than you should

If you inject more vosoritide than you should, contact your child's doctor straight away.

If you forget to use vosoritide

If your child misses a dose, the injection should still be given if it is within 12 hours of the scheduled time. If more than 12 hours have passed since the scheduled dose time, do not inject the missed dose. Wait until the next day and continue with the usual dose at the usual time.

If you stop using vosoritide

Always talk to your child's doctor before deciding to stop your child's treatment. If you have any further questions on the use of this medicine, ask your child's doctor, nurse or pharmacist.

Please see accompanying package leaflet.

Travelling with vosoritide

Short- and long-term travel requires considerable planning and organisation. Bring enough treatment supply and injection materials with you in case of unforeseen circumstances, e.g. if you decide to stay longer than planned, or if a needle breaks, etc. Keep the treatment reliably cool. Make sure to give injections to your child at a suitable time.

How to store vosoritide

Keep this medicine **out of the sight and reach of children. Do not freeze.** Store in a refrigerator (2°C – 8°C) in the original package in order to protect from light. Vosoritide may be stored at room temperature (below 30°C) for **up to 90 days**, but not beyond the expiry date. **Record** on the carton **the date** you remove vosoritide from the refrigerator to store at room temperature. **Do not** return vosoritide to refrigerator after storage at room temperature.

Reporting of side effects

If your child gets any side effects, talk to your child's doctor, nurse or pharmacist. This includes any possible side effects not listed in the package leaflet. By reporting side effects you can help provide more information on the safety of this medicine.