

IMPORTANT INFORMATION ABOUT COMBINED HORMONAL CONTRACEPTIVES AND RISK OF BLOOD CLOTS

All combined hormonal contraceptives (CHCs) increase the risk of having a blood clot. The overall risk of a blood clot due to a CHC is small but clots can be serious and may in very rare cases even be fatal.

It is very important that you recognise when you might be at greater risk of a blood clot, what signs and symptoms you need to look out for and what action you need to take.

In which situations is the risk of a blood clot highest?

- in the first year of using a CHC (including if you are re-starting use after a break of 4 weeks or more) - if you are very overweight
- if you are older than 35 years
- if you have a family member who has had a blood clot at a relatively young age (eg below 50) - if you have given birth in the previous few weeks

If you smoke and are over 35 years old you are strongly advised to stop smoking or use a non-hormonal method of contraception.

Seek medical attention immediately if you experience any of the following symptoms:

- Severe pain or swelling in either of your legs that may be accompanied by tenderness, warmth or changes in the skin colour such as turning pale, red or blue. You may be experiencing a **deep vein thrombosis**.
- Sudden unexplained breathlessness or rapid breathing; severe pain in the chest which may increase with deep breathing; sudden cough without an obvious cause (which may bring up blood);. You may be experiencing a serious complication of deep vein thrombosis called a **pulmonary embolism**. This occurs if the blood clot travels from the leg to the lung.
- Chest pain, often acute, but sometimes just discomfort, pressure, heaviness, upper body discomfort radiating to the back, jaw, throat, arm together with a feeling of fullness associated with indigestion or choking, sweating, nausea, vomiting or dizziness. You may be experiencing a **heart attack**
- Face, arm or leg weakness or numbness, especially on one side of the body; trouble speaking or understanding; sudden confusion; sudden loss of vision or blurred vision; severe headache/migraine that is worse than normal. You may be experiencing a **stroke**.

Watch out for symptoms of a blood clot, especially if you have: •

Just had an operation

- been off your feet for a long time (eg. because of an injury or illness, or if your leg is in a cast) • a long journey (more than about 4 hours)

Remember to tell your doctor, nurse or surgeon that you are taking a CHC if you: •

Are due to or have had surgery

- Are in any situation when a healthcare professional asks you if you are taking any medications

Suspected Adverse Drug Reactions (side effects) or medication errors may be reported using the Medicines Authority ADR reporting form, which is available online at <http://www.medicinesauthority.gov.mt/adrportal>, and sent by post or email to; P: Pharmacovigilance Section at Post-Licensing Directorate, Medicines Authority, Sir Temi Żammit Buildings, Malta Life Sciences Park, San Ġwann SĠN 3000 E: postlicensing.medicinesauthority@gov.mt or alternatively to JV Healthcare Limited P: JV Healthcare Limited, Navi Buildings, Pantar Road, Lija LJA 2021 E: info@jvpharma.eu