

# **REBLOZYL® (luspatercept)**

## **Prescriber's Checklist**

Important information for healthcare providers prescribing REBLOZYL for women of childbearing potential

The Prescriber's Checklist is to be used before initiating treatment, at each administration, and then at regular intervals when performing follow-up.

For more information about REBLOZYL or to obtain further copies of this document, please contact AM Mangion Ltd on 00 n356 2397 6333

Please see the REBLOZYL SmPC for complete prescribing information.

REBLOZYL® (luspatercept) Prescriber's Checklist

REBLOZYL

Prescriber's Checklist for Women of Childbearing Potential

| Patient Identification | Prescriber Details |
|------------------------|--------------------|
| Name:                  | Name:              |
|                        | Signature:         |
|                        | Date:              |

| Prior to Initiation of Treatment   |
|--|
| Treatment with REBLOZYL is contraindicated in pregnancy and in women of childbearing potential not using at least one highly effective method of contraception <ul style="list-style-type: none"> <li>- There are no data from the use of REBLOZYL in pregnant women. Studies in animals have shown reproductive toxicity and embryo-foetal toxicity. Clinical implications are potential foetal loss and teratogenicity.</li> </ul> |
| <input type="checkbox"/> Provide counselling before treatment initiation regarding the potential teratogenic risk of REBLOZYL and required actions to minimise this risk.  |
| <input type="checkbox"/> Inform women of childbearing potential of the necessity for at least one highly effective method of contraception while on treatment and for 3 months after discontinuation.  |
| <input type="checkbox"/> A pregnancy test must be carried out and a negative result must be verified in women of childbearing potential before starting treatment.   |
| <input type="checkbox"/> Provide the Patient Card to a women of childbearing potential.  |
| Duration of Treatment  |
| <input type="checkbox"/> Provide regularly counselling regarding the potential teratogenic risk of REBLOZYL and required actions to minimise this risk   |
| <input type="checkbox"/> Remind women of childbearing potential that they must use at least one highly effective method of contraception during treatment with REBLOZYL.   |
| During treatment with REBLOZYL, women must not become pregnant. If a woman becomes pregnant or wants to become pregnant, REBLOZYL should be discontinued.  |
| During treatment with REBLOZYL, pregnancy tests must be repeated at suitable intervals and medically verified as negative.   |
| Discontinuation of Treatment   |
| <input type="checkbox"/> Counsel women of childbearing potential that at least one highly effective method of contraception should be maintained for at least 3 months following discontinuation of treatment with REBLOZYL.   |
| <input type="checkbox"/> Provide counselling in the event of pregnancy and evaluation of the outcome of any pregnancy. <ul style="list-style-type: none"> <li><input type="checkbox"/> Not applicable (this patient did not become pregnant while on treatment or within 3 months of discontinuation of REBLOZYL.)</li> </ul>  |

**Should a pregnancy occur during treatment or within 3 months following discontinuation of treatment with REBLOZYL, remind the patient that it should be reported to the prescriber, Medicines Authority and to BMS by AM Mangion Ltd on Tel 00 356 23976333 and email - pv@ammangion.com , irrespective of adverse outcomes observed.**

***This Prescriber checklist fulfils the conditions of the marketing authorisation and has been approved by the competent authority, version number 2007-MT-2500001 and date of approval by the NCA "<month> <year>".***

## REPORTING OF ADVERSE REACTIONS

Suspected adverse reactions and medication errors should be reported at  
ADR Reporting, The Medicines Authority, Post-Licensing Directorate,  
Sir Temi Zammit Buildings, Malta Life Sciences Park,  
San Gwann SGN 3000, Malta  
Website: [www.medicinesauthority.gov.mt](http://www.medicinesauthority.gov.mt)  
e-mail: [postlicensing.medicinesauthority@gov.mt](mailto:postlicensing.medicinesauthority@gov.mt)

OR

ADR Reporting: [www.medicinesauthority.gov.mt/adrportal](http://www.medicinesauthority.gov.mt/adrportal)

AND

AM Mangion Ltd  
Mangion House  
New Street off Valletta Road  
Luqa LQA6000, Malta  
Email: [pv@ammangion.com](mailto:pv@ammangion.com)

Tel – 00 356 23976333